

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/101264/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Pritchard, Manon F. , Powell, Lydia C. , Jack, Alison A., Powell, Kate , Beck, Konrad , Florance, Hannah, Forton, Julian, Rye, Philip D., Dessen, Arne, Hill, Katja E. and Thomas, David W. 2017. A low-molecular-weight alginate oligosaccharide disrupts pseudomonal microcolony formation and enhances antibiotic effectiveness. *Antimicrobial Agents and Chemotherapy* 61 (9) , e00762-17. 10.1128/AAC.00762-17

Publishers page: <http://dx.doi.org/10.1128/AAC.00762-17>

Please note:


Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



1 **FIGURES**


2



3

4 **FIG S1** Circular dichroism (CD) spectra of high Mw pseudomonal alginate mixed with OligoG CF-5/20. Scans 1-7 show spectra of high Mw
5 alginate (~20 μ M) followed over ~77 min upon heating from 4 to 37 $^{\circ}$ C; scans 8-11 (~44min) are recorded after addition of OligoG CF-5/20 (850
6 μ M) followed by addition of Ca $^{2+}$ (1mM) (scans 12-17, ~ 66min). Spectra of OligoG CF-5/20 (850 μ M) alone and its sum with high Mw alginate
7 (20 μ M) are shown as black and red dashed lines, respectively.


8



10

11 **FIG S2** CD spectra of high Mw pseudomonal alginate mixed with OligoG CF-5/20. Scans 1-2 show spectra of high Mw alginate (~20 μM)
 12 followed over ~20 min after heating from 4 to 37°C, after which Ca^{2+} was added to a final concentration of 1 mM (scans 3-8) and 2 mM (scans
 13 9-11). Oligo G CF-5/20 was added to $c_{\text{fin}} = 700\mu\text{M}$ (scans 12-15; molar ratio high Mw alginate to OligoG CF-5/20 1:50) and $c_{\text{fin}} = 4.2\text{mM}$ (scans
 14 16-22; molar ratio high Mw alginate to OligoG CF-5/20 1:600. Adding calcium to $c_{\text{fin}} = 9\text{mM}$ results in spontaneous alginate precipitation (scans
 15 23-25).

16



17

18 **FIG S3** Biofilm disruption assay showing SEM images of established (24 h) *P. aeruginosa* (A) PAO1 and (B) NH57388A biofilms treated for 24
 19 h with 2% OligoG CF-5/20 ± colistin (16 µg/ml). Scale bar 20 µm.

20