

# Quality-of-life outcomes from the Prostate Adenocarcinoma: TransCutaneous Hormones (PATCH) trial evaluating luteinising hormone-releasing hormone agonists versus transdermal oestradiol for androgen suppression in advanced prostate cancer

Duncan C. Gilbert\*, Trinh Duong\*, Howard G. Kynaston†, Abdulla A. Alhasso‡, Fay H. Cafferty\*, Stuart D. Rosen§, Subramanian Kanaga-Sundaram¶, Sanjay Dixit\*\*\*, Marc Laniado††, Sanjeev Madaan†††, Gerald Collins††††, Alvan Pope†††, Andrew Welland\*, Matthew Nankivell\*, Richard Wassersug\*\*\*, Mahesh K. B. Parmar\*, Ruth E. Langley\* and Paul D. Abel†††††††

\*Medical Research Council Clinical Trials Unit at University College London, London, †Cardiff School of Medicine, Cardiff University, Cardiff, ‡The Beatson West of Scotland Cancer Centre, Glasgow, §National Heart and Lung Institute, Imperial College London, London, ¶Mid-Yorkshire Hospitals NHS Trust, Pinderfields General Hospital, Wakefield, \*\*\*Scunthorpe General Hospital, North Lincolnshire and Goole NHS Trust, Scunthorpe, ††Frimley Health NHS Foundation Trust, Wexham Park Hospital, Slough, †††Dartford and Gravesham NHS Trust, Darent Valley Hospital, Dartford, §§Stockport NHS Foundation Trust, Stepping Hill Hospital, Stockport, ††††The Hillingdon Hospitals NHS Foundation Trust, London, UK, \*\*\*University of British Columbia, Vancouver, BC, Canada, †††††Imperial College Healthcare NHS Trust, and †††††††Imperial College London, London, UK

D.C.G., T.D., R.E.L. and P.D.A. equal contribution.

## Objectives

To compare quality-of-life (QoL) outcomes at 6 months between men with advanced prostate cancer receiving either transdermal oestradiol (tE2) or luteinising hormone-releasing hormone agonists (LHRHa) for androgen-deprivation therapy (ADT).

## Patients and methods

Men with locally advanced or metastatic prostate cancer participating in an ongoing randomised, multicentre UK trial comparing tE2 versus LHRHa for ADT were enrolled into a QoL sub-study. tE2 was delivered via three or four transcutaneous patches containing oestradiol 100 µg/24 h. LHRHa was administered as per local practice. Patients completed questionnaires based on the European Organisation for Research and Treatment of Cancer quality of life questionnaire 30-item core (EORTC QLQ-C30) with prostate-specific module QLQ PR25. The primary outcome measure was global QoL score at 6 months, compared between randomised arms.

## Results

In all, 727 men were enrolled between August 2007 and October 2015 (412 tE2, 315 LHRHa) with QoL questionnaires completed at both baseline and 6 months. Baseline clinical characteristics

were similar between arms: median (interquartile range) age of 74 (68–79) years and PSA level of 44 (19–119) ng/mL, and 40% (294/727) had metastatic disease. At 6 months, patients on tE2 reported higher global QoL than those on LHRHa (mean difference +4.2, 95% confidence interval 1.2–7.1;  $P = 0.006$ ), less fatigue, and improved physical function. Men in the tE2 arm were less likely to experience hot flushes (8% vs 46%), and report a lack of sexual interest (59% vs 74%) and sexual activity, but had higher rates of significant gynaecomastia (37% vs 5%). The higher incidence of hot flushes among LHRHa patients appear to account for both the reduced global QoL and increased fatigue in the LHRHa arm compared to the tE2 arm.

## Conclusion

Patients receiving tE2 for ADT had better 6-month self-reported QoL outcomes compared to those on LHRHa, but increased likelihood of gynaecomastia. The ongoing trial will evaluate clinical efficacy and longer term QoL. These findings are also potentially relevant for short-term neoadjuvant ADT.

## Keywords

androgen-deprivation therapy, quality of life, transdermal oestradiol, #ProstateCancer, #PCSM

## Introduction

Prostate cancer is the most frequent cancer diagnosis in men in the developed world and responsible for 11 000 deaths per year in the UK and 26 000 in the USA [1,2]. Prostate cancer cell growth is driven by androgen signalling, and androgen-deprivation therapy (ADT) forms a cornerstone of treatment. Evidence supports the use of ADT in conjunction with radiotherapy in localised [3,4] and locally-advanced disease [5,6], and as first-line therapy in the metastatic setting [7].

ADT, usually achieved using LHRH agonists (LHRHa) in contemporary practice, is associated with numerous side-effects [8,9]. Specifically, these include declining bone health [10,11], weight gain and metabolic syndrome [12], sexual dysfunction [13–15], hot flushes [16,17], mental and cognitive decline [18–22], and physical deterioration and fatigue [23–26]. LHRHa increase the risk of depression in men with prostate cancer [14], reportedly driven by the loss of sexual function [27]. Recent data suggest an increased risk of subsequent Alzheimer's disease [28]. An association with increased cardiac events is described but remains controversial [29]. Whereas a number of interventions have been shown to ameliorate the toxicities of LHRHa to a greater or lesser extent [8], further efforts are required to maintain the highest possible quality of life (QoL) for these patients.

PATCH (Prostate Adenocarcinoma: TransCutaneous Hormones, MRC PR09) is an ongoing randomised controlled trial comparing transdermal oestradiol (tE2) delivered via transcutaneous patches vs LHRHa in men with advanced prostate cancer. LHRHa act through the hypothalamic-pituitary axis to suppress testosterone production by the testes. Endogenous E2 in men is derived from testosterone through aromatase. Thus, it is also suppressed by, and consequently contributes to, the toxicity profile of LHRHa [9]. Exogenous administration of E2 inhibits the hypothalamic-pituitary axis (thereby suppressing testosterone) as well but maintains E2 levels and hence mitigates some of the toxicity of LHRHa. Administration of exogenous E2 via oral or i.v. routes is associated with risk of thrombosis and adverse cardiovascular events [30]. However, tE2 avoids the hepatic first-pass effects mediating these risks, as supported by previous results from PATCH (254 patients) showing similar rates of cardiovascular events in both tE2 and LHRHa arms after a median follow-up of 19 months [31]. Among this initial cohort, castration rates were similar in both arms.

In the present report, we compare QoL outcomes at 6 months from randomisation between the two hormonal treatments, based on data available from ~700 patients.

## Patients and Methods

The study design for the PATCH trial has previously been described [31]. Briefly, patients from participating UK centres were eligible for recruitment if they had locally advanced or metastatic prostate cancer, and a treatment plan for indefinite ADT in the metastatic setting or ≥3 years for locally advanced disease. National regulatory and ethics committees approved the protocol, and participating hospitals obtained the appropriate local approvals. Participants provided written informed consent.

Men were randomly allocated (in a 2:1 ratio before February 2011 and then 1:1) to receive tE2 or LHRHa (open-label). This was done centrally according to a computer-based minimisation algorithm with a random element (80%), balanced for the following factors: disease stage, age, smoking status, personal or family history of heart disease, which LHRHa agent was to be used, PSA level, intent to give radical radiotherapy, and centre.

Patients in the tE2 arm received, after a dose regimen change in August 2007 [32], four FemSeven patches (oestradiol 100 µg/24 h), which were self-administered and changed twice weekly during the first 4 weeks. This was reduced to three patches changed twice weekly, provided testosterone levels were <1.7 nmol/L. LHRHa was administered as per local practice.

## QoL Data Collection

Patients received a specific patient information sheet for the QoL study and provided separate consent to participate in this component of the study. QoL information was collected on paper questionnaires using the European Organisation for Research and Treatment of Cancer quality of life questionnaire 30-item core (EORTC QLQ-C30) and the prostate-specific module QLQ-PR25. These were self-completed by participants, who were instructed to record responses without discussion with site staff, friends or relatives. Data were collected before randomisation, then at 4, 8 and 12 weeks, and subsequently every 3 months up to 2 years after randomisation. The QLQ-C30 includes a range of domains that are either multi-item scales or single-item measures: a global health status/QoL scale, five function scales (physical, role, emotional, cognitive, and social), three symptom scales (fatigue, pain, and nausea and vomiting), and six single items (dyspnoea, insomnia, appetite loss, constipation, diarrhoea, and financial difficulties). QLQ-PR25 contains 25 items designed to assess QoL in patients with prostate cancer, including urinary, bowel and sexual symptoms and functioning, and hormone-related symptoms.

The Independent Data Monitoring Committee (IDMC) permitted release of the QoL data during the first 6 months

from randomisation for patients already enrolled whilst the main trial continues.

### Statistical Analysis

For each multi-item QoL domain (e.g. global QoL), a summary score was derived according to the EORTC QLQ-C30 scoring manual [33], with a range of 0–100. For example, the summary global QoL score is a standardised average of the patients' scores from the questions 'How would you rate your overall health during the past week?' and 'How would you rate your overall quality of life?'. These scores were considered as continuous variables. A higher score corresponds to improved outcomes for global QoL and function scales, but indicates more symptoms (hence poorer outcomes) for symptoms scales. Single-item domains (e.g. sexual interest) were analysed based on reported responses on the questionnaires ('not at all', 'a little', 'quite a bit', or 'very much').

The primary outcome was global QoL score at 6 months, as differences in hormone-related symptoms potentially impacting on QoL were expected to be apparent by then [17,20,25,26,31]. The following domains were secondary outcomes: sexual interest, sexual activity, whether feeling less masculine as result of illness or treatment, cognitive functioning, physical functioning, fatigue, and selected hormone-related symptoms of hot flushes, gynaecomastia, and weight gain. Gynaecomastia was reported as sore or enlarged nipples or breasts.

Patients were considered to have baseline QoL data, if they completed their first QoL questionnaire either by the date of randomisation or 1 week after, but before starting trial treatment. Information on QoL outcomes at 6 months was based on the questionnaire completed nearest to this time point, within a ± 3-month window.

Multi-item QoL domains at 6 months were compared between randomised arms using Tobit regression models (to account for scores being bounded by 0 and 100) [34], adjusting for baseline score. Single-item domains were categorised according to pre-defined binary outcomes for comparison between arms (for ease of clinical interpretation); for example, hot flushes were analysed as 'quite a bit'/'very much' vs 'not at all'/'a little'. These were compared between arms using logistic regression models, adjusting for baseline response.

All models were further adjusted for the following pre-defined baseline factors: age, calendar year (partly to account for the change in allocation ratio), smoking status, stage of disease (M0/M1), and whether patient was newly diagnosed or relapsing. All comparisons between arms were based on the original allocated treatment, and included patients randomised after the change in patch dose regimen [32] who

had data on the relevant QoL domains at both baseline and 6 months. A significance level of 0.05 was chosen a priori, without adjustment for multiple statistical testing. Additional exploratory analyses were undertaken to investigate associations between global QoL and other domains.

Statistical analyses were performed using Stata version 14 (Stata Corporation, College Station, TX, USA).

### Results

Between 14 August 2007 and 5 October 2015, 875 men were recruited under the revised patch dose regime, 480 allocated to tE2 and 395 to LHRHa. Within the tE2 arm, 468 patients enrolled on the QoL sub-study, of whom 412 (86% of 480) completed QoL questionnaires at both baseline and 6 months. For the LHRHa group, 385 participated in the QoL sub-study, with 315 (80% of 395) having both baseline and 6-month QoL data available (Fig. 1). Baseline clinical characteristics were similar between arms for the 727 patients included in the 6-month QoL analyses (Table 1). The overall median (interquartile range) age was 74 (68–79) years and PSA level was 44 (19–119) ng/mL, and 40% (294/727) had metastatic disease. There were no differences in baseline global QoL by age or testosterone level, but men with T4 tumours had worse global QoL compared with other T-stages, and patients with metastatic disease had worse baseline QoL than M0 patients.


Rates of castration were equivalent between the LHRHa and tE2 arms at both 3 and 6 months; the proportion of patients with testosterone concentrations ≤1.7 nmol/L was 93.6% for LHRHa and 93.7% for tE2 at 3 months, and 89.8% and 92.2% at 6 months, respectively.

At 6 months, global QoL declined from baseline in both arms (Table 2), but to a lesser extent in the tE2 patients (mean change -2.8) compared to those on LHRHa (-5.0). The estimated mean difference in 6-month global QoL between arms was +4.2 (95% CI 1.2–7.1;  $P = 0.006$ ) in favour of tE2. There was no evidence that the treatment effect on global QoL at 6 months differed by age ( $\leq 70$  vs  $> 70$  years; test for interaction  $P = 0.56$ ).

In addition, there was less decline in physical function among tE2 patients (mean change -2.8 vs -5.7), with a mean difference in 6-month score of +5.8 (95% CI 2.8–8.8;  $P < 0.001$ ) between arms. In addition, tE2 patients had less fatigue at 6 months, mean difference between arms -4.3 (95% CI -8.1 to -0.6;  $P = 0.02$ ) favouring the patches. However, there was no difference in reported decline in cognitive function between arms.

Analysis of specific domains linked with testosterone suppression (Table 3) showed that tE2 patients were less likely than LHRHa patients to report having no interest in sex [59% vs 74%; odds ratio (OR) 0.42, 95% CI 0.28–0.62;

**Figure 1** Consolidated Standards of Reporting Trials (CONSORT) diagram of patient inclusion in the analysis of 6-month QoL data. <sup>a</sup>The allocation ratio was 1:2 LHRHa:tE2 before 21/02/2011 and 1:1 thereafter. <sup>b</sup>Patients were considered to have baseline QoL data if they had completed their first QoL questionnaire either by the date of randomisation or within 1 week after but before starting trial treatment. <sup>c</sup>As overall survival is a co-primary outcome measure in the ongoing trial, the number of patients who have died before completing 6-month QoL questionnaire is not provided.



$P < 0.001]$  and being ‘not at all’ sexually active (78% vs 87%; OR 0.51, 95% CI 0.32–0.82;  $P = 0.005$ ). Interestingly, there was weak evidence that the negative effect of LHRHa compared to tE2 on interest in sex was more pronounced in patients aged  $\leq 70$  years than those  $> 70$  years ( $t$ -test for interaction  $P = 0.06$ ).

The likelihood of experiencing ‘quite a bit’ or ‘very much’ hot flushes was significantly lower in the tE2 group (8% vs 46%; OR 0.10, 95% CI 0.07–0.16;  $P < 0.001$ ). However, as expected, patients in the tE2 arm were much more likely to report ‘quite a bit’ or ‘very much’ gynaecomastia than those receiving LHRHa (37% vs 5%; OR 12.70, 95% CI

7.14–22.60;  $P < 0.001$ ). There was no difference between arms in patients who reported feeling ‘quite a bit’ or ‘very much less’ masculine (as a result of their illness or treatment) or experiencing ‘quite a bit’ or ‘very much’ weight gain.

An association between hot flushes and deterioration in global QoL was seen in both arms at 6 months, with patients who experienced more severe symptoms reporting lower scores (Table 4,  $P < 0.001$ ). The relationship between gynaecomastia and global QoL was assessed in the tE2 arm only, owing to few LHRHa patients reporting symptoms. Gynaecomastia was associated with poorer global QoL at

**Table 1** Patient characteristics for those with both baseline and 6-month QoL questionnaires completed (727 patients).

| Variable  | LHRHa       | tE2         |
|---|-------------|-------------|
| Number of patients  | 315         | 412         |
| Age (years) at randomisation                                |             |             |
| <70, n (%)  | 99 (31)     | 128 (31)    |
| 70–79, n (%)  | 146 (46)    | 202 (49)    |
| ≥80, n (%)  | 70 (22)     | 82 (20)     |
| Median (IQR)  | 74 (67–79)  | 73 (68–79)  |
| Metastatic disease, n (%)                                   | 133 (42)    | 161 (39)    |
| Bone metastases (% of those with metastatic disease), n (%) | 120 (90)    | 148 (92)    |
| PSA concentration (ng/mL)                                   |             |             |
| <50, n (%)  | 173 (55)    | 214 (52)    |
| 50–<500, n (%)  | 121 (39)    | 163 (40)    |
| ≥500, n (%)   | 19 (6)      | 35 (9)      |
| Median (IQR)  | 43 (22–115) | 45 (18–119) |
| Tumour stage, n (%)   |             |             |
| T0/1/2  | 19 (6)      | 27 (7)      |
| T3  | 220 (70)    | 296 (72)    |
| T4  | 52 (17)     | 64 (16)     |
| TX  | 24 (8)      | 25 (6)      |
| N category, n (%)   |             |             |
| N0  | 118 (37)    | 145 (35)    |
| N+  | 87 (28)     | 102 (25)    |
| NX  | 110 (35)    | 165 (40)    |
| Gleason sum score, n (%)                                    |             |             |
| 4–6   | 28 (10)     | 28 (8)      |
| 7   | 89 (31)     | 137 (37)    |
| 8–10  | 171 (59)    | 207 (56)    |
| Smoking status, n (%)                                       |             |             |
| Never smoked  | 119 (38)    | 167 (41)    |
| Previous smoker   | 162 (51)    | 204 (50)    |
| Current smoker  | 34 (11)     | 41 (10)     |
| WHO performance status, n (%)                               |             |             |
| Normal activity   | 209 (66)    | 293 (71)    |
| Avoid strenuous activity                                    | 92 (29)     | 102 (25)    |
| Up and about >50%   | 14 (4)      | 17 (4)      |
| Year of randomisation, n (%)                                |             |             |
| 2007/2008   | 30 (10)     | 69 (17)     |
| 2009/2010   | 40 (13)     | 75 (18)     |
| 2011/2012   | 141 (45)    | 156 (38)    |
| 2013/2015   | 104 (33)    | 112 (27)    |

IQR, interquartile range.

6 months (Table 5,  $P = 0.004$ ), although the adverse effect was only seen in patients reporting ‘very much’ gynaecomastia (corresponding to 8% of the group with data available). Other QoL domains associated with lower global QoL score were: poorer cognitive and physical function, increased fatigue, weight gain, and feeling less masculine (data not shown).

After accounting for hot flushes, there was little difference in the 6-month global QoL score between arms (estimated mean difference comparing tE2 vs LHRHa  $-0.4$ , 95% CI  $-3.8$  to  $3.0$ ;  $P = 0.80$ ). In comparison, the difference between arms remained after other QoL domains were individually adjusted for (data not shown). This suggests a significant component of the effect of treatment arm on global QoL could potentially be attributable to the higher incidence of hot flushes in LHRHa patients.

In addition, there was an association between severity of hot flushes and fatigue at 6 months in both arms (data not shown), which may potentially account for the increased fatigue in the LHRHa vs tE2 arm; after adjusting for hot flushes, there was little difference in the 6-month fatigue score between arms (mean difference comparing tE2 vs LHRHa  $0.0$ , 95% CI  $-4.3$  to  $4.4$ ;  $P = 0.98$ ). Further *post hoc* analyses showed a relationship between hot flushes and sleep disturbance within both arms; 72% (124/172) of patients reporting ‘quite a bit’ or ‘very much’ hot flushes had trouble sleeping compared to 43% (232/534) of those with ‘not at all’ or ‘a little’ hot flushes ( $P < 0.001$ , with similar results by arm).

Patients experiencing gynaecomastia were more likely to report feeling less masculine at 6 months, with 24% (36/148) of men who reported ‘quite a bit’ or ‘very much’ gynaecomastia feeling ‘quite a bit’ or ‘very much’ less masculine compared to 7% (17/247) of those reporting ‘not at all’ or ‘a little’ gynaecomastia ( $P < 0.001$ ). The protocol explicitly allowed prophylactic breast bud radiotherapy and 5% of patients on tE2 received this treatment as opposed to no patients on LHRHa. Two patients underwent surgical treatment for gynaecomastia who were both on tE2, corresponding to 0.4% (2/480) of the overall tE2 arm cohort enrolled to date.

## Discussion

In the present study, we found better overall QoL after 6 months of ADT with tE2 compared to LHRHa, as well as less fatigue and improved physical function. While the magnitude of the QoL effects was modest [35], some additional differences are important to note. Men treated with LHRHa were more likely to report lack of sexual interest (74% vs 59%) and being not sexually active (87% vs 78%) [20]. In addition, tE2 patients had lower rates of hot flushes but more gynaecomastia, consistent with earlier findings from the trial [31].

Significant hot flushes were reported by 8% of men on tE2 compared to 44% of those on LHRHa. Interestingly, there was a suggestion that hot flushes mediated the treatment effect on global QoL, potentially accounting for both the reduced global QoL and increased fatigue in the LHRHa compared to tE2 arm. Conversely, 37% men on tE2 reported significant gynaecomastia compared with 5% on LHRHa, although gynaecomastia was only seen to adversely affect global QoL if the patient reported ‘very much’ symptoms (which corresponded to <10% of the tE2 cohort). It is noteworthy that men may vary significantly in how bothersome gynaecomastia is on an individual basis [36]. In addition, data from the main PATCH trial suggest no association between E2 levels and clinical gynaecomastia (data not shown).

**Table 2** QoL multi-item domains: scores at 6 months by treatment arm.\*

| Outcome            | Arm   | Number of patients | Mean score at baseline (95% CI) | Mean score at 6 months (95% CI) | Mean change in 6-month score from baseline (95% CI) | Mean difference in 6-month score between arms (95% CI) | P comparing arms |
|--------------------|-------|--------------------|---------------------------------|---------------------------------|---|--|------------------|
| Global QoL score   | LHRHa | 308                | 75.1 (72.7, 77.4)               | 70.1 (67.7, 72.4)               | -5.0 (-7.4, -2.7)                                   |  |                  |
|                    | tE2   | 403                | 78.0 (76.1, 80.0)               | 75.2 (73.3, 77.2)               | -2.8 (-4.7, -0.8)                                   | +4.2 (1.2, 7.1)  | 0.006            |
| Cognitive function | LHRHa | 309                | 86.9 (84.8, 89.0)               | 82.8 (80.7, 84.9)               | -4.1 (-6.2, -2.0)                                   |  |                  |
|                    | tE2   | 403                | 87.5 (85.7, 89.3)               | 84.0 (82.2, 85.9)               | -3.5 (-5.3, -1.6)                                   | +1.9 (-1.8, 5.5)                                       | 0.32             |
| Physical function  | LHRHa | 307                | 87.6 (85.4, 89.8)               | 81.8 (79.6, 84.1)               | -5.7 (-7.9, -3.5)                                   |  |                  |
|                    | tE2   | 399                | 89.0 (87.2, 90.9)               | 86.2 (84.3, 88.1)               | -2.8 (-4.7, -1.0)                                   | +5.8 (2.8, 8.8)  | <0.001           |
| Fatigue            | LHRHa | 304                | 18.9 (16.4, 21.4)               | 27.2 (24.7, 29.7)               | 8.3 (5.8, 10.8)                                     |  |                  |
|                    | tE2   | 400                | 17.1 (14.8, 19.4)               | 23.0 (20.8, 25.3)               | 6.0 (3.7, 8.2)                                      | -4.3 (-8.1, -0.6)                                      | 0.02             |

\*For global QoL, cognitive function and physical function, a higher score corresponds to a better outcome. For fatigue, a higher score corresponds to more fatigue.

**Table 3** QoL single-item domains: proportion of patients with pre-defined outcomes at 6 months by treatment arm.

| Outcome                                   | Arm   | Number of patients | Number (%) of patients with outcome at baseline | Number (%) of patients with outcome at 6 months | OR (95% CI)         | P comparing arms |
|---|-------|--------------------|---|---|---------------------|------------------|
| Felt quite a bit/very much less masculine | LHRHa | 292                | 16 (5)  | 41 (14)   |                     |                  |
|   | tE2   | 381                | 18 (5)  | 53 (14)   | 0.92 (0.57, 1.48)   | 0.73             |
| Not at all interested in sex              | LHRHa | 268                | 118 (44)  | 199 (74)  |                     |                  |
|   | tE2   | 350                | 155 (44)  | 208 (59)  | 0.42 (0.28, 0.62)   | <0.001           |
| Not at all sexually active                | LHRHa | 266                | 168 (63)  | 231 (87)  |                     |                  |
|   | tE2   | 348                | 214 (61)  | 271 (78)  | 0.51 (0.32, 0.82)   | 0.005            |
| Quite a bit/very much gynaecomastia       | LHRHa | 293                | 4 (1)   | 14 (5)  |                     |                  |
|   | tE2   | 386                | 2 (1)   | 144 (37)  | 12.70 (7.14, 22.60) | <0.001           |
| Quite a bit/very much hot flushes         | LHRHa | 291                | 6 (2)   | 135 (46)  |                     |                  |
|   | tE2   | 390                | 9 (2)   | 32 (8)  | 0.10 (0.07, 0.16)   | <0.001           |
| Quite a bit/very much weight gain         | LHRHa | 296                | 13 (4)  | 21 (7)  |                     |                  |
|   | tE2   | 379                | 13 (3)  | 27 (7)  | 1.06 (0.56, 2.00)   | 0.87             |

LHRHa therapy can severely impact on physical well-being and other QoL outcomes [14–16]. Hot flushes, reported by 40–80% of men on LHRHa [17,37,38], are linked to sleep disturbance and psychological distress [16,38]. In our present study, patients with hot flushes had more trouble sleeping, which may account for the effect of hot flushes on increased fatigue and reduced QoL. tE2 appeared to be effective in reducing the severity of hot flushes in men on ADT in a prior study, consistent with our present findings [39]. The adverse effects of LHRHa on sexual outcomes, which can have significant psychological impact on both patients and their partners, have also been well-documented [13–15]. Data from men castrated for reasons other than prostate cancer suggest exogenous oestrogen can help maintain sexual interest [40,41]. Other potential benefits of tE2 reported include protective effects on cognition [42], although we did not find a difference in cognitive function between arms within our present study, possibly because short-term outcomes were analysed and/or limitations of the questionnaires used for assessing the cognitive domain.

Several strategies have been investigated in an attempt to mitigate the adverse effects LHRHa therapy [8]. Randomised trials have shown some benefit of medoxyprogesterone, venlafaxine and gabapentin in reducing hot flushes associated with LHRHa, and exercise may improve levels of fatigue and overall QoL [43–45]. Agents which can potentially preserve bone health during treatment with LHRHa include bisphosphonates, denosumab or toremifene [8]. Importantly, however, data from PATCH recently showed that patients on tE2 avoid the loss in bone mineral density seen with LHRHa administration [46]. The data presented here suggest tE2 as an alternative to LHRHa might limit the requirement for additional treatments to allay the side-effects of LHRHa over and above bone health. Alternatively a low dose of tE2 in addition to LHRHa could be investigated in the future as a treatment for bothersome hot flushes.

Alternatively, an intermittent approach to ADT has been assessed for clinical efficacy and potential QoL benefits. In the non-metastatic setting, intermittent ADT appears not to be

**Table 4** Global QoL score at 6 months in both treatment arms, by patients' experience of hot flushes.

| Experienced hot flushes at 6 months | LHRHa arm                       |                                 |   | tE2 arm                         |                                 |   | Mean difference in 6-month score (95% CI), both arms combined* |
|-------------------------------------|---------------------------------|---------------------------------|---|---------------------------------|---------------------------------|---|--|
|                                     | Number of patients (% of total) | Mean score at 6 months (95% CI) | Mean change in 6-month score from baseline (95% CI) | Number of patients (% of total) | Mean score at 6 months (95% CI) | Mean change in 6-month score from baseline (95% CI) |  |
| Not at all                          | 77 (26)                         | 75.1 (70.4, 79.8)               | -0.9 (-5.6, 3.9)                                    | 288 (73)                        | 78.6 (76.4, 80.8)               | -1.8 (-4.0, 0.4)                                    | Reference group  |
| A little                            | 83 (28)                         | 74.0 (69.9, 78.1)               | -4.0 (-8.1, 0.1)                                    | 75 (19)                         | 65.1 (60.6, 69.6)               | -5.9 (-10.4, -1.4)                                  | -6.8 (-10.6, -3.1)   |
| Quite a bit                         | 92 (31)                         | 65.6 (61.1, 70.1)               | -8.1 (-12.6, -3.5)                                  | 23 (6)                          | 67.4 (58.0, 76.7)               | -5.8 (-15.1, 3.6)                                   | -10.4 (-15.0, -5.9)  |
| Very much                           | 47 (16)                         | 62.4 (56.6, 68.2)               | -7.6 (-13.4, -1.9)                                  | 10 (3)                          | 65.0 (52.7, 77.4)               | -2.5 (-14.8, 9.9)                                   | -11.8 (-17.6, -6.0)  |
|                                     |                                 |                                 |   |                                 |                                 |   | P < 0.001  |

\*Estimated from Tobit regression models, adjusted for treatment arms, baseline global QoL score and other pre-defined baseline factors. There was no evidence that the effect of hot flushes on 6-month global QoL score differed by treatment arm (P for interaction = 0.20).

**Table 5** Global QoL score at 6 months in the tE2 arm\*, by whether patient reported to have experienced gynaecomastia.

| Experienced sore or enlarged nipples or breasts | Number of patients (% of total) | Mean score at baseline (95% CI) | Mean score at 6 months (95% CI) | Mean change in 6-month score from baseline (95% CI) | Mean difference in 6-month score† (95% CI) |
|---|---------------------------------|---------------------------------|---------------------------------|---|--|
| Not at all                                      | 55 (14)                         | 78.6 (73.0, 84.3)               | 73.3 (67.9, 78.7)               | -5.3 (-10.7, 0.1)                                   | Reference group                            |
| A little  | 190 (49)                        | 80.0 (77.2, 82.7)               | 78.0 (75.2, 80.9)               | -1.9 (-4.8, 0.9)                                    | 4.8 (-0.8, 10.4)                           |
| Quite a bit                                     | 114 (29)                        | 74.4 (70.9, 78.0)               | 73.8 (70.5, 77.0)               | -0.7 (-3.9, 2.6)                                    | 3.3 (-2.7, 9.3)                            |
| Very much                                       | 32 (8)                          | 74.5 (66.7, 82.2)               | 63.8 (56.1, 71.5)               | -10.7 (-18.4, -3.0)                                 | -7.6 (-15.6, 0.4)                          |
|   |                                 |                                 |                                 |   | P = 0.004                                  |

\*This association was not assessed in the LHRHa arm owing to few patients reporting symptoms. †Estimated from Tobit regression models, adjusted for baseline global QoL score and other pre-defined baseline factors.

inferior to continuous therapy in terms of overall survival, with some potential benefits as regards hot flushes, libido and possibly fatigue, but not global health [47]. However, a randomised trial by Hussain et al. [48] including 1535 men with metastatic prostate cancer failed to show non-inferiority for intermittent ADT based on overall survival. Although small improvements were initially seen for sexual function and mental health, older men assigned to intermittent ADT had no apparent reduction in bone, endocrine, or cognitive events and experienced an increased incidence of ischaemic and thrombotic events [49].

It is increasingly apparent across a number of QoL domains that there are important differences in the unintended consequences of ADT depending upon the method chosen to achieve castrate levels of testosterone [50,51]. Here, we have shown that at 6 months of treatment, tE2 improves patients' QoL in a number of domains compared to LHRHa, i.e. fewer hot flushes, less fatigue, improved physical functioning, sexual interest and sexual activity, but at a cost of increased incidences of gynaecomastia. This can be viewed in addition to the beneficial effects on tE2 on bone mineral density previously reported within PATCH [46], also noting the lack of any excess cardiovascular or thromboembolic effects from tE2 [31]. From our present data, hot flushes appear to potentially account for the increased fatigue and reduced global QoL among patients on LHRHa.

We acknowledge the relatively short-term outcomes assessed and presented here. However, ADT is often used for periods as short as 6 months when administered as neoadjuvant therapy along with radiotherapy to treat localised disease. As such, our present 6-month QoL data are clinically pertinent, given short-term neoadjuvant ADT has been shown to be associated with impaired QoL [52]. Further data from the ongoing trial will inform whether the differences between arms persist long term. Although it is premature to suggest a fundamental change in practice when it comes to starting patients on ADT, comprehensive analysis of comparative efficacy and toxicity within PATCH will allow men and their partners to optimise treatment choices.

## Acknowledgements

We thank the National Institute for Health Research (NIHR) Cancer Research Network for staff support; all the patients who participated in the PATCH trial and their families; the research staff at the participating hospitals; the PATCH Trial Management Group, Trial Steering Committee, and the Independent Data Monitoring Committee (see Appendix for list of members). We also thank all previous and current members of the PATCH trial team including: Gordana Jovic who currently maintains the statistical database for the trial; Anna Bara for overseeing the practical running of the trial; Robin Carpenter for his assistance with the data management;

and Ben Spittle who was the PATCH trial manager during 2011–2014.

## Conflict of Interests

Ruth E. Langley has served as an advisor for, and received honoraria from Bayer. Alvan J. Pope receives meeting sponsorship from Ipsen Ltd and is a shareholder in AstraZeneca. The other authors declare they have no conflicts of interest.

## Funding/support and Role of the Sponsor

The PATCH study is funded by Cancer Research UK, grant number C17093/A12443 (trial CRUK/06/001) and University College London (UCL), and is sponsored by Imperial College London. The funding sources and sponsor had no role in the study design; collection, management, analysis, or interpretation of the data; preparation, review, or approval of the manuscript; nor decision to submit the manuscript for publication.

## References

- 1 National Cancer Institute. SEER Stat Fact Sheets: Prostate Cancer. Available at: <http://seer.cancer.gov/statfacts/html/prost.html>. Accessed June 2016.
- 2 Cancer Research UK. Prostate cancer statistics. Available at: <http://wwwcancerresearchukorg/health-professional/cancer-statistics/statistics-by-cancer-type/prostate-cancer>. Accessed June 2016.
- 3 D'Amico AV, Chen MH, Renshaw AA, Loffredo M, Kantoff PW. Androgen suppression and radiation vs radiation alone for prostate cancer: a randomized trial. *JAMA* 2008; 299: 289–95
- 4 Jones CU, Hunt D, McGowan DG, et al. Radiotherapy and short-term androgen deprivation for localized prostate cancer. *N Engl J Med* 2011; 365: 107–18
- 5 Bolla M, de Reijke TM, Van Tienhoven G, et al. Duration of androgen suppression in the treatment of prostate cancer. *N Engl J Med* 2009; 360: 2516–27
- 6 Warde P, Mason M, Ding K, et al. Combined androgen deprivation therapy and radiation therapy for locally advanced prostate cancer: a randomised, phase 3 trial. *Lancet* 2011; 378: 2104–11
- 7 James ND, Spears MR, Clarke NW, et al. Survival with newly diagnosed metastatic prostate cancer in the “docetaxel era”: data from 917 patients in the control arm of the STAMPEDE trial (MRC PR08, CRUK/06/019). *Eur Urol* 2015; 67: 1028–38
- 8 Nguyen PL, Alibhai SM, Basaria S, et al. Adverse effects of androgen deprivation therapy and strategies to mitigate them. *Eur Urol* 2014; 67: 825–36
- 9 Phillips I, Shah S, Duong T, Abel P, Langley RE. Androgen deprivation therapy and the re-emergence of parenteral oestrogen in prostate cancer. *Oncol Hematol Rev* 2014; 10: 42–7
- 10 Diamond TH, Higano CS, Smith MR, Guise TA, Singer FR. Osteoporosis in men with prostate carcinoma receiving androgen-deprivation therapy: recommendations for diagnosis and therapies. *Cancer* 2004; 100: 892–9
- 11 Shahinian VB, Kuo YF, Freeman JL, Goodwin JS. Risk of fracture after androgen deprivation for prostate cancer. *N Engl J Med* 2005; 352: 154–64
- 12 Saylor PJ, Smith MR. Metabolic complications of androgen deprivation therapy for prostate cancer. *J Urol* 2009; 181: 1998–2008
- 13 Benedict C, Traeger L, Dahn JR, et al. Sexual bother in men with advanced prostate cancer undergoing androgen deprivation therapy. *J Sex Med* 2014; 11: 2571–80
- 14 Donovan KA, Walker LM, Wassersug RJ, Thompson LM, Robinson JW. Psychological effects of androgen-deprivation therapy on men with prostate cancer and their partners. *Cancer* 2015; 121: 4286–99
- 15 Higano CS. Sexuality and intimacy after definitive treatment and subsequent androgen deprivation therapy for prostate cancer. *J Clin Oncol* 2012; 30: 3720–5
- 16 Savard J, Hervouet S, Ivers H. Prostate cancer treatments and their side effects are associated with increased insomnia. *Psychooncology* 2013; 22: 1381–8
- 17 Gonzalez BD, Jim HS, Donovan KA, et al. Course and moderators of hot flash interference during androgen deprivation therapy for prostate cancer: a matched comparison. *J Urol* 2015; 194: 690–5
- 18 Cary KC, Singla N, Cowan JE, Carroll PR, Cooperberg MR. Impact of androgen deprivation therapy on mental and emotional well-being in men with prostate cancer: analysis from the CaPSURE registry. *J Urol* 2014; 191: 964–70
- 19 McGinty HL, Phillips KM, Jim HS, et al. Cognitive functioning in men receiving androgen deprivation therapy for prostate cancer: a systematic review and meta-analysis. *Support Care Cancer* 2014; 22: 2271–80
- 20 Lee M, Jim HS, Fishman M, et al. Depressive symptomatology in men receiving androgen deprivation therapy for prostate cancer: a controlled comparison. *Psychooncology* 2015; 24: 472–7
- 21 Gonzalez BD, Jim HS, Booth-Jones M, et al. Course and predictors of cognitive function in patients with prostate cancer receiving androgen-deprivation therapy: a controlled comparison. *J Clin Oncol* 2015; 33: 2021–7
- 22 Dinh KT, Reznor G, Muralidhar V, et al. Association of androgen deprivation therapy with depression in localized prostate cancer. *J Clin Oncol* 2016; 34: 1905–12
- 23 Alibhai SM, Breunis H, Timilshina N, et al. Long-term impact of androgen-deprivation therapy on physical function and quality of life. *Cancer* 2015; 121: 2350–7
- 24 Sadetsky N, Greene K, Cooperberg MR, Hubbard A, Carroll PR, Satariano W. Impact of androgen deprivation on physical well-being in patients with prostate cancer: analysis from the CaPSURE (Cancer of the Prostate Strategic Urologic Research Endeavor) registry. *Cancer* 2011; 117: 4406–13
- 25 Gonzalez BD, Jim HS, Small BJ, et al. Changes in physical functioning and muscle strength in men receiving androgen deprivation therapy for prostate cancer: a controlled comparison. *Support Care Cancer* 2016; 24: 2201–7
- 26 Nelson AM, Gonzalez BD, Jim HS, et al. Characteristics and predictors of fatigue among men receiving androgen deprivation therapy for prostate cancer: a controlled comparison. *Support Care Cancer* 2016; 24: 4159–66
- 27 Sharpley CF, Christie DR, Bitsika V, Miller BJ. Trajectories of total depression and depressive symptoms in prostate cancer patients receiving six months of hormone therapy. *Psychooncology* 2016. [Epub ahead of print]. doi: 10.1002/pon.4100.
- 28 Nead KT, Gaskin G, Chester C, et al. Androgen deprivation therapy and future Alzheimer's disease risk. *J Clin Oncol* 2016; 34: 566–71
- 29 Nguyen PL, D'Amico AV. Toward personalizing the use of androgen deprivation therapy to maximize benefit and minimize harm. *Eur Urol* 2015; 68: 397–8
- 30 Byar DP. Proceedings: The Veterans Administration Cooperative Urological Research Group's studies of cancer of the prostate. *Cancer* 1973; 32: 1126–30
- 31 Langley RE, Cafferty FH, Alhasso AA, et al. Cardiovascular outcomes in patients with locally advanced and metastatic prostate cancer treated with luteinising-hormone-releasing-hormone agonists or transdermal oestrogen:

- the randomised, phase 2 MRC PATCH trial (PR09). *Lancet Oncol* 2013; 14: 306–16
- 32 Langley RE, Godsland IF, Kynaston H, et al. Early hormonal data from a multicentre phase II trial using transdermal oestrogen patches as first-line hormonal therapy in patients with locally advanced or metastatic prostate cancer. *BJU Int* 2008; 102: 442–5
- 33 Aaronson NK, Ahmedzai S, Bergman B, et al. The European Organization for Research and Treatment of Cancer QLQ-C30: a quality-of-life instrument for use in international clinical trials in oncology. *J Natl Cancer Inst* 1993; 85: 365–76
- 34 Austin PC, Escobar M, Kopec JA. The use of the Tobit model for analyzing measures of health status. *Qual Life Res* 2000; 9: 901–10
- 35 Cocks K, King MT, Velikova G, Martyn St-James M, Fayers PM, Brown JM. Evidence-based guidelines for determination of sample size and interpretation of the European Organisation for the Research and Treatment of Cancer Quality of Life Questionnaire Core 30. *J Clin Oncol* 2011; 29: 89–96.
- 36 Wassersug RJ, Oliffe JL. The social context for psychological distress from iatrogenic gynecomastia with suggestions for its management. *J Sex Med* 2009; 6: 989–1000
- 37 Walker LM, Tran S, Robinson JW. Luteinizing hormone-releasing hormone agonists: a quick reference for prevalence rates of potential adverse effects. *Clin Genitourin Cancer* 2013; 11: 375–84
- 38 Ulloa EW, Salup R, Patterson SG, Jacobsen PB. Relationship between hot flashes and distress in men receiving androgen deprivation therapy for prostate cancer. *Psychooncology* 2009; 18: 598–605
- 39 Gerber GS, Zagaja GP, Ray PS, Rukstalis DB. Transdermal estrogen in the treatment of hot flushes in men with prostate cancer. *Urology* 2000; 55: 97–101
- 40 Wibowo E, Wassersug RJ. The effect of estrogen on the sexual interest of castrated males: Implications to prostate cancer patients on androgen-deprivation therapy. *Crit Rev Oncol Hematol* 2013; 87: 224–38
- 41 Handy AB, Jackowich RA, Wibowo E, Johnson TW, Wassersug RJ. Gender preference in the sexual attractions, fantasies, and relationships of voluntarily castrated men. *Sex Med* 2016; 4: e51–9
- 42 Engler-Chiurazzi EB, Brown CM, Povroznik JM, Simpkins JW. Estrogens as neuroprotectants: Estrogenic actions in the context of cognitive aging and brain injury. *Prog Neurobiol* 2016. [Epub ahead of print]. doi: 10.1016/j.pneurobio.2015.12.008.
- 43 Bourke L, Gilbert S, Hooper R, et al. Lifestyle changes for improving disease-specific quality of life in sedentary men on long-term androgen-deprivation therapy for advanced prostate cancer: a randomised controlled trial. *Eur Urol* 2014; 65: 865–72
- 44 Cormie P, Galvao DA, Spry N, et al. Can supervised exercise prevent treatment toxicity in patients with prostate cancer initiating androgen-deprivation therapy: a randomised controlled trial. *BJU Int* 2015; 115: 256–66
- 45 Bourke L, Smith D, Steed L, et al. Exercise for men with prostate cancer: a systematic review and meta-analysis. *Eur Urol* 2016; 69: 693–703
- 46 Langley RE, Kynaston HG, Alhasso AA, et al. A Randomised comparison evaluating changes in bone mineral density in advanced prostate cancer: luteinising hormone-releasing hormone agonists versus transdermal oestradiol. *Eur Urol* 2016; 69: 1016–25
- 47 Crook JM, O'Callaghan CJ, Duncan G, et al. Intermittent androgen suppression for rising PSA level after radiotherapy. *N Engl J Med* 2012; 367: 895–903
- 48 Hussain M, Tangen CM, Berry DL, et al. Intermittent versus continuous androgen deprivation in prostate cancer. *N Engl J Med* 2013; 368: 1314–25
- 49 Herszman DL, Unger JM, Wright JD, et al. Adverse health events following intermittent and continuous androgen deprivation in patients with metastatic prostate cancer. *JAMA Oncol* 2016; 2: 453–61
- 50 Sun M, Choueiri TK, Hamnvik OP, et al. Comparison of gonadotropin-releasing hormone agonists and orchectomy: effects of androgen-deprivation therapy. *JAMA Oncol* 2016; 2: 500–7
- 51 Albertsen PC, Klotz L, Tombal B, Grady J, Olesen TK, Nilsson J. Cardiovascular morbidity associated with gonadotropin releasing hormone agonists and an antagonist. *Eur Urol* 2014; 65: 565–73
- 52 Gay HA, Michalski JM, Hamstra DA, et al. Neoadjuvant androgen deprivation therapy leads to immediate impairment of vitality/hormonal and sexual quality of life: results of a multicenter prospective study. *Urology* 2013; 82: 1363–8

**Correspondence:** Duncan C. Gilbert, MRC Clinical Trials Unit at UCL, Institute of Clinical Trials and Methodology, Aviation House, 125 Kingsway, London WC2B 6NH, UK.

**e-mail:** duncan.gilbert@ucl.ac.uk

**Abbreviations:** ADT, androgen-deprivation therapy; EORTC QLQ-C30, European Organisation for Research and Treatment of Cancer quality of life questionnaire 30-item core; LHRHa, LHRH agonists; OR, odds ratio; PATCH, Prostate Adenocarcinoma: TransCutaneous Hormones; QoL, quality of life; (t)E2, (transdermal) oestradiol.

## Appendix

### PATCH trial committees

Trial Management Group (in alphabetical order): Paul Abel (Chief Investigator), Abdulla A. Alhasso, Anna Bara, Robin Carpenter, Noel W. Clarke, David Dearnaley, Trinh Duong, Duncan Gilbert, Ian F. Godsland, Gordana Jovic, Ruth E. Langley, Howard G. Kynaston, Roger Kockelbergh, Mahesh K.B. Parmar, Michael Philips (patient representative), Stuart D. Rosen, Andrew Welland.

Trial Steering Committee: David Guthrie (Chair), John Chester, and Richard Cowan.

Independent Data Monitoring Committee: Peter Hoskin (Chair), Philip Smith, and Laurence Collette.