

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <http://orca.cf.ac.uk/105884/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Harun, A, Finlay, Andrew, Salek, M. Sam and Piguet, Vincent 2018. How to train to discharge a dermatology outpatient: A review. *Dermatology* 233 (4) , pp. 260-267. 10.1159/000479060 file

Publishers page: <http://dx.doi.org/10.1159/000479060> <<http://dx.doi.org/10.1159/000479060>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



8th June 2017

“Clean” manuscript, with revisions

TITLE PAGE

Title

How to train to discharge a dermatology outpatient: a review

Authors

N. A. Harun, A. Y. Finlay, M. S. Salek, V. Piguat

Name of institution at which the work was conducted

Department of Dermatology and Wound Healing,

Division of Infection and Immunity,

School of Medicine, Cardiff University

3rd Floor, Glamorgan House

Heath Park, Cardiff CF14 4XN, UK

Running head

How to discharge an outpatient

Name, address, telephone, fax and email of corresponding author

Prof A Y Finlay,

Department of Dermatology and Wound Healing

Division of Infection and Immunity,

School of Medicine, Cardiff University

3rd Floor, Glamorgan House

Heath Park, Cardiff CF14 4XN, UK

Tel No: +44 (0) 7860 112639

Fax No: +44 (0) 29 2074 4312

Email: FinlayAY@cf.ac.uk

Names and address of other authors:

Dr N A Harun,

Department of Dermatology and Wound Healing

Division of Infection and Immunity,

School of Medicine, Cardiff University

3rd Floor, Glamorgan House

Heath Park, Cardiff CF14 4XN, UK

Tel No: +44(0)29 2074 4721

Fax No: +44(0)29 2074 4312

Email: ainitaharun@gmail.com

Department of Dermatology,
University Malaya Medical Centre,
Kuala Lumpur 50603, Malaysia

Prof S Salek

School of Life and Medical Sciences
University of Hertfordshire, College Lane,
Hatfield AL10 9AB, UK

Email: mssalek@googlemail.com

Prof V Piguet

Department of Dermatology and Wound Healing
Division of Infection and Immunity,
School of Medicine, Cardiff University
3rd Floor, Glamorgan House
Heath Park, Cardiff CF14 4XN, UK

Tel No: +44 (0) 29 2074 4721

Fax No: +44 (0) 29 2074 4312

Email: Piguenv@cf.ac.uk

ORCID numbers

N A Harun: 0000 0002 8272 9123

A Y Finlay: 0000 0003 2143 1646

S Salek: 0000 0002 4612 5699

V Piguet: 0000 0001 6079 4517

Key words

Education, discharge decision, dermatology outpatients, clinical practice

Key Message

This review describes how to educate trainees to make high quality appropriate outpatient discharge decisions.

Manuscript word, Table, Figure and Reference count

Abstract: 229 (max 250)

Word count: 2,450

Tables: 5

Figures: 1

References: 20

Conflict of interest

All authors state no conflict of interest.

MANUSCRIPT

Abstract

Background/Aims

The decision to discharge is a critical and common outpatient consultation event. However little guidance exists over how discharge decision taking can be taught. We aimed to provide educational recommendations concerning outpatient discharge decision taking.

Methods

Recommendations were drawn from prior interviews with 40 consultant dermatologists and 56 dermatology outpatients, and from the 'Traffic-light' design discharge information checklist, developed using the Delphi technique.

Results

The key strategies to follow to appropriately manage the outpatient discharge process are: to warn patients in advance, to understand patients' agendas, to allow extra time for the discharge process, to prepare patients to self-manage, to provide a 'safety-net' and provide the GP with a clear management plan. Aspects to be considered include patient mobility, presence of carer, type of employment, diagnostic certainty and use of checklist or guidelines. Key training aspects include teaching structured thought processes when discharging, discharging according to context, developing communication and negotiation skills, avoiding decision biases and encouraging good

inter-professional collaboration. Training should include the consideration of the possibility of discharge each consultation.

Novel training strategies have been developed on how to appropriately manage the outpatient discharge process, including involving and informing patients. These strategies focus on safe decision-making, being patient-centred and organizing an efficient healthcare service framework.

Conclusion

Structured outpatient discharge training for dermatologists is now possible, based on information from detailed doctor and patient based qualitative studies.

Introduction

The decision whether or not to discharge an outpatient has major implications for quality of patient care and efficiency of outpatient services [1-5]. This is particularly important in the primarily outpatient specialty of dermatology, where very large numbers of patients are seen and delayed discharge of patients has a major impact on service efficiency. Both patients and the health service can be adversely affected by inappropriately early outpatient discharge, and conversely by inappropriate follow-up appointments.

Most clinicians have never been taught how to discharge an outpatient and most trainees receive no discharge training [2]. Despite outpatient discharge being one of the commonest decisions taken by clinicians there has been very little research aimed at understanding it. Training on how to discharge is essential to ensure that appropriate decisions are made in the best interests of the patient. Training is also necessary because of long waiting lists for outpatient specialist care [6,7], uncertainties over how

to make discharge decisions [8,9], patients' lack of understanding of discharge instructions [10,11], mis-matched perception during discharge discussions [4,10] and because of delays in re-accessing secondary care [4].

Discharge decision-making is a specific decision skill encompassing clinical expertise, communication skills, weighing of ethical issues and ability to organise clinical practice. It needs to be taught to improve the quality of care and to reduce medical errors. Education research has focused on the most effective ways to teach clinicians how to make appropriate diagnostic and treatment decisions [12], but discharge decision-making has largely been ignored. What little has been published has focused on educating nurses, patients and family members [11] with little emphasis on educating clinicians. Vreeland et al. suggested [11] that inpatient discharge education should be structured, individualised, include patient's preferences, include family involvement, provide multimedia instructions, emphasise medication intake, and include education of the multidisciplinary team involved in inpatient discharge: clinicians, nurses, dieticians and social workers. They concluded that discharge education reduced re-admissions and lowered healthcare costs [11].

We recently reviewed the literature [2], reported dermatologists' views on influences on the outpatient discharge decision [3] and patients' views on being discharged [4]. We proposed a "Traffic-light" design dermatology outpatient discharge information checklist to ensure that the minimum information to take a quality discharge decision is available to the clinician [5]. During this qualitative research project consultants and patients made many suggestions over the discharge process and about key aspects that could be taught to trainee doctors.

The aim of this paper is to provide recommendations, bringing together lessons from these prior studies that are of direct relevance to the education of clinicians concerning outpatient discharge decision-making. This paper does not present novel research

findings, but encapsulates the educational and training issues arising from recent research [2-5].

Background

The educational recommendations given in this paper are based on information gathered in three qualitative studies of consultant and patient perspectives of dermatology outpatient discharge [3-5]. Ethical permission was given for the gathering of educational information during these studies. Forty dermatologists working in 11 hospitals were interviewed [3]: they were asked on how they arrived at a discharge or follow-up decision, what critical factors influenced their decision, what they considered an “appropriate” discharge and were asked to suggest strategies on how junior clinicians should be taught to take appropriate discharge decisions. Fifty-six patients with a range of dermatology conditions were interviewed after a routine dermatology outpatient consultation at which a decision was reached about whether or not they should be discharged [4]. In order to identify the most important pieces of information that a clinician should know to take an appropriate outpatient discharge decision [5], 17 dermatologists took part in a three-stage Delphi process, that led to the creation of a Traffic-light design checklist [5].

Educational Recommendations

Methods to improve the process of outpatient discharge

Key methods to improve the process of outpatient discharge, based on the three qualitative studies [3-5], are given in table 1. Key aspects of an appropriate discharge decision are given in fig. 1.

Methods to inform the discharge decision to ensure its appropriateness

The 13 items (table 2) of the “Traffic-light” design discharge checklist [5] can be used as a teaching tool. In the Delphi process that led to this checklist some items did not

qualify as part of the discharge checklist, but are still useful to consider in specific clinical contexts during the discharge decision-making process (table 3). During the studies, consultants suggested several methods on how dermatology trainees could be trained to take discharge decisions. Table 4 summarises these methods, illustrated with relevant consultant quotes.

Proposed educational objectives

Since outpatient discharge is the critical endpoint of patient care in the outpatient clinical setting, appropriate discharge decisions are not only important for the well-being of individual patients, but are integral to ensuring healthcare service efficiency. The purpose of educating clinicians about outpatient discharge is to enable them to make high quality dermatology outpatient discharge decisions, with the aim of ensuring that no patient is disadvantaged as a result of a clinician's lack of confidence and competence in taking discharge decisions. We propose a series of broad educational objectives (table 5).

Discussion

We have developed novel strategies on how to appropriately manage the outpatient discharge process, inform and make discharge decisions. There are three main aspects: safer decision-making (self-awareness, knowledge on limits of own expertise, patients' characteristics, diagnostic certainty, providing contact after discharge, clear treatment plan), patient-centred decisions (accurate perception and management of patients' expectations, reducing clinician biases, longer consultation time for 'difficult' patients, effective communication) and increased service efficiency (tool to guide discharge decisions, use of clinical guidelines, good time management, effective collaboration

with colleagues and general practitioners, provision of open appointments and discharge policies). These strategies are similar to those identified in a qualitative study of discharge from general medical outpatients [10].

Factors of most importance to patients in being satisfied with their dermatology outpatient consultation have been identified as physician's concern about and focus on quality of life, physician's interest in patients' symptoms and medical problems and the carrying out of full skin examination or specialised examination such as use of a dermatoscope [14]. That study however did not investigate satisfaction with the discharge process. In a study of dermatology outpatients immediately after being discharged, we identified considerable dissatisfaction amongst some patients at being discharged, even though they had apparently agreed to the discharge during the consultation [4].

There is a distinction between what actions are critical to the process of making an appropriate discharge and how best to inform patients about the discharge decision itself. Both these aspects of patient discharge involve a set of decision skills that could be taught in a structured manner [2-5].

The consideration of a wide array of clinical and nonclinical factors influencing the discharge process can make judgement difficult for the inexperienced clinician who may inadvertently be subject to inappropriate biases, often without insight [2,3]. Junior doctors should be educated on the likely existence of bias, in order to support better clinical decision-taking [12,15].

In many health care systems there is great pressure on clinical time, with restricted consultation length. It may also take longer to discharge a patient than to offer a follow-up appointment. However unnecessary follow-up appointments may be a major contributing factor to the time pressures on dermatology outpatient clinics as every unnecessary appointment is a lost opportunity to see a new patient or to spend longer

time with other patients. Training in outpatient discharge decision taking may therefore contribute appropriately to increased service efficiency. Most clinicians have never been trained to take outpatient discharge decisions but have learnt themselves. Some clinicians may have little insight into the influences on their decision taking and little awareness of the biases that they bring to clinical decision taking. This may contribute to scepticism of the possible benefits of training in discharge decision taking.

One limitation of the information on which the educational recommendations are made is that the information came primarily from detailed interviews with consultant dermatologists and with patients, but without input from general practitioners, other health care professionals or from dermatology trainees. Further research gaining the perspective of these other groups [16] would be of benefit. The research on which these educational recommendations were based was carried out in the context of the UK publically funded National Health Service. While many of the influences on dermatology outpatient discharge decisions may be encountered in any healthcare system, it is possible that there may be other influences and different strategies to be considered in, for example, a private dermatology clinic. We are not aware of any research on this topic carried out in a private health care setting.

There are many different patterns of private clinical practice both within and between different countries, with widely differing rates of outpatient discharge. In private practice, the dynamics between patient and physician may differ from public practice, with the additional influence of consultation fees. However even in private practice it may be helpful for the physician to have received training in clinical strategies relating to taking appropriate discharge decisions and in managing the process of discharge to the satisfaction of the patient. It remains to be tested whether the concepts and strategies we have described may be appropriate for use in private practice.

One educational objective is to encourage clinicians to consider the items in the “Traffic-light” design discharge information checklist [5] and to use the checklist routinely in outpatient practice. The checklist covers five disease-related items, four patient empowerment items and four patient concerns, which if considered, may encourage earlier appropriate discharge or prevent inappropriately early discharge. The five disease-related items are in the red (stop) section of the checklist. These indicate a command to the clinician to stop and seriously consider these items before proceeding. When these have been considered, the clinician proceeds to the next step, considering the four patient empowerment items in the yellow (proceed with caution) section. After this, the clinician should consider addressing concerns items in the green (permission to proceed) section. When all of these items have been addressed, the patient can be discharged [5]. Although it was designed to be concise and user friendly, the use of the checklist as a practical tool in the clinic or in education has yet to be tested. The use of the checklist may be the method most likely to ascertain discharge appropriateness and its use may help to minimise biases due to personal judgement and cognitive errors, which recur predictably in discharging patients. We suggest prospective research to ascertain whether the use of this discharge information checklist and education in this area can make any difference to the quality of outpatient discharge decision making. It is possible that both the discharge information checklist and suggested educational goals and strategies may be used as a platform to improve discharge practice in other specialties or also in inpatient discharge practice.

When first encountered, the Traffic-light checklist approach may seem unfeasible. However in a small survey in a publicly funded dermatology clinic [5]. 61% of 18 clinicians reported that they would like to use the Traffic light checklist. The Traffic light concept has not been used in private practice. Prospective research is required to

test whether an adaptation of the concept or perhaps a different approach may be needed for the private practice setting.

There is not yet sufficient evidence to support the Traffic light system being a requirement for training. However training concerning discharge decision taking should be a requirement, as this is one of the commonest clinical decisions that every dermatologist takes on a daily basis. Whether or not it is used in clinical practice, the Traffic light checklist may be useful to focus discussion and training in this area.

A radically different alternative approach to reducing outpatient clinic numbers by increasing discharge decision rates has been suggested [17]. A senior clinician reviewed all outpatient charts two weeks before the clinic and a brief written management plan was made. This resulted in a large reduction in clinic sizes. It would be possible to use such a review as a teaching exercise for trainees.

It is important to encourage trainees to reflect [18] on how well they handle judgement under uncertainty, how effectively they communicate and to have insight into their unconscious use of mental shortcuts that may cause biased decisions. Over a century ago, the Flexner Report [19] stressed that medical students should be taught how to apply clinical reasoning when making decisions when faced with uncertainty, and the discharge decision is a good example on which to base education about this issue.

Clinicians are often immersed in the technicalities of disease management and therefore may neglect the more “humane” aspects of patient care. They may fail to realise that during consultations patients are affected by the attitudes of their clinician [4,10,20]. Teaching trainee clinicians to actively reflect over each of their patient consultations may assist trainees to project an effortless empathic response to patients. Such skills are learned partly from experience and need frequent self-reflection to maintain and improve them. Because of the often incurable nature and high visibility of skin diseases,

trainee dermatologists need to be more aware of the needs of patients with skin disease and whether a patient is actually ready for discharge.

Other aspects of routine clinical decision making such as diagnostic and treatment decisions should also be taught, rather than assuming that clinicians will learn by imperceptibly “soaking up” the experience of seniors in an apprentice relationship. These wider decisions should be examined in a similar way, to help transform the “Art of Medicine” into an approach less liable to bias. It is appropriate to try to improve the quality of clinician’s approach to taking clinical decisions, since clinicians exert great influence on their patients and the healthcare team. It is also important for those who make decisions about groups of patients or populations to learn a more structured and transparent approach to making evidence-based decisions. Integrating training in decision making into dermatology trainee programmes will hopefully improve the quality of patient care.

Conclusion

This paper summarizes a novel educational approach to the process of making outpatient discharge decisions, based on qualitative research evidence. The time has come to revisit and sharpen our skills over discharging outpatients in an attempt to minimise biases and uncertainties. When genuine effort, professional wisdom and patient empowerment come together, the goal of making more appropriate discharge decisions may then be realised. If dermatologists better understood how discharge decisions should be carried out, greater discharge decision appropriateness might be achieved in the best interests of patients and dermatology services. Furthermore, it is hoped that educators in medicine will seriously consider inclusion of “outpatient discharge decision-making” in the mainstream curriculum.

Acknowledgements

We wish to thank the dermatologists and dermatology patients whose ideas are summarised in this article. We also wish to thank clinicians from the University Hospital Wales (UHW), Cardiff who contributed suggestions on making outpatient discharge decisions at meetings organised by AYF in 2008-2009, incorporated in table 1.

References

- 1 Sullivan FM. How do clinicians decide to discharge someone from their outpatient clinic? *J Manag Med* 1993; 7: 24-8.
- 2 Harun NA, Salek S, Piguet V, Finlay AY. The dermatology outpatient discharge decision: understanding a critical but neglected process. *Br J Dermatol* 2014; 170: 1029-38.
- 3 Harun NA, Finlay AY, Salek MS, Piguet V. Appropriate and inappropriate influences on outpatient discharge decision making in dermatology: a prospective qualitative study. *Br J Dermatol* 2015; 173: 720-30.
- 4 Harun NA, Finlay AY, Piguet V, Salek S. Understanding clinician influences and patient perspectives on outpatient discharge decisions: a qualitative study. *BMJ Open* 2017; 7(3): e010807.
- 5 Harun N, Finlay A, Salek MS, Piguet V. The development and clinical evaluation of a “Traffic-light design” dermatology outpatient discharge information checklist. *Br J Dermatol* 2016; 175: 572-82.

- 6 Salek MS, Siyani S, Basra MK, Finlay AY. The clinical and non-clinical factors influencing discharge decisions in dermatology: Is there a need for discharge strategy? *Int J Clin Pharm* 2012; 34 (1):178-179.
- 7 Stainkey LA, Seidl IA, Johnson AJ et al. The challenge of long waiting lists: how we implemented a GP referral system for non-urgent specialist' appointments at an Australian public hospital. *BMC Health Serv Res* 2010; 10: 1.
- 8 Burkey Y, Black M, Reeve H et al. Long-term follow-up in outpatient clinics. 2: The view from the specialist clinic. *Fam Pract* 1997; 14: 29-33.
- 9 Wu K-H, Chen IC, Li C-J et al. The influence of physician seniority on disparities of admit/discharge decision making for ED patients. *Am J Emerg Med* 2012; 30: 1555-60.
- 10 Burkey Y, Black M, Reeve H. Patients' views on their discharge from follow up in outpatient clinics: qualitative study. *BMJ* 1997; 315: 1138-41.
- 11 Vreeland D, Rea RE, Montgomery L. A Review of the Literature on Heart Failure and Discharge Education. *Crit Care Nurs Q* 2011; 34: 235-45.
- 12 Hunink MM, Weinstein MC, Wittenberg E et al. *Decision making in health and medicine: integrating evidence and values*: Cambridge University Press. 2014.
- 13 Brahams D. The medicolegal implications of teleconsulting in the UK. *J Telemed Telecare* 1995; 1: 196-201.
- 14 De Salins CA, Brenaut E, Misery L, Roguedas-Contios AM. Factors influencing patient satisfaction: assessment in outpatients in dermatology department. *J Eur Acad Dermatol Venereol* 2016; 30: 1823-1828.
- 15 Croskerry P, Norman G. Overconfidence in clinical decision making. *The Am J Med* 2008; 121: S24-S9.

- 16 Reeve H, Baxter K, Newton P, Burkey Y, Black M, Roland M. Long term follow-up in outpatient clinics.1: The view from general practice. Family Practice 1997; 14: 24-28.
- 17 Donnellan F, Hussain T, Aftab AR, McGurk C. Reducing unnecessary outpatient attendances. Int J Health Care Quality Assurance 2010; 23: 527-531.
- 18 Magnussen, L Trotter C.M. The reflective clinician. Nurse Educator 1997; 22(3): 40-44.
- 19 Flexner, A. 1910. Medical education in the United States and Canada [Online].
http://archive.carnegiefoundation.org/pdfs/elibrary/Carnegie_Flexner_Report.pdf [Accessed: 16th April 2017].
- 20 Agledahl KM, Gulbrandsen P, Førde R et al. Courteous but not curious: how doctors' politeness masks their existential neglect. A qualitative study of video-recorded patient consultations. J Med Ethics 2011; 37: 650-4.

Table 1. Key methods for discharging outpatients, based on three qualitative studies [3,4,5].

Key methods for outpatient discharge
1. Have discharge in mind in all consultations.
2. At first appointment set expectations for discharge. <i>Explain the likely course of treatment and expected outcomes (e.g. acne or laser patients may have predictable number of clinic appointments).</i>
3. Discuss possible discharge “at the next visit” and document it in patient’s notes.
4. Ask patient directly if they want to come back to the clinic. <i>Be firm and directive where appropriate.</i>

5. Explore the patient's agenda.

Ask patient what they were hoping for from the consultation.

Take more time when handling a "difficult" patient.

6. Be honest if you have nothing more to offer the patient.

7. Allow extra time to explain reasons and plan the discharge with the patient.

Time is the key to a successful discharge.

Clinic numbers may need to be reduced, but compensated by a higher rate of discharge.

8. Inform and summarise events and investigation results in a positive manner.

Remind patients of the positive outcomes from their clinic visits.

Remind patients that we don't want them to have unnecessary clinic visits.

9. Prepare patients to self-manage.

Educate patients and provide information about realistic outcomes

Provide information to help patients self-manage e.g. information leaflets or website links.

Give positive feedback, e.g. praising the patient for learning about their disease and taking more responsibility for management.

10. Provide patients with a safety net.

Make clear that care is not ending but being transferred back to the GP.

Encourage patients to seek help from their GP and explain the circumstances when the GP would re-refer them back to the clinic..

Use email to communicate with patients after 'simple' surgical procedures e.g. uncomplicated basal cell carcinoma in low risk site. Note that emailing patients may have medicolegal implications [13].

Allow direct patient contact with the consultant's secretary, especially with anxious patients.

If there is a dermatology department specialist nurse advice service, explain to patients when and how to access this service

Consider giving the patient an 'open access' appointment.

11. Provide GP with a clear management plan. Consider copying this 'discharge' letter to patient.

Check that the GP can prescribe your recommended therapy or carry out your

suggested plan.

Dictate the discharge letter in front of the patient.

Table 2. The 13 items of the ‘Traffic-light’ design discharge information checklist [5].

Disease related	Patient empowerment	Addressing concerns
Am I certain of the diagnosis?	Has the patient understood the diagnosis and treatment outcome?	Has the patient any concerns?
Is the condition severe?	Have I explained to the patient a clear plan of treatment?	Can the patient re-access secondary care easily if the problem recurs?
Is the patient on the appropriate treatment?	Have I explained the treatment side effects?	Is the patient happy to be discharged?
Can the patient be managed in primary care?	Has the patient understood how to self-manage?	Am I happy to discharge the patient?
Will the patient benefit from my follow up?		

Table 3. Other items to be considered when making a discharge decision, in addition to those in the ‘Traffic-light’ checklist.

Items to be considered	Clinical context
Patient’s psychological state of mind	Patients who are anxious
Complexity of the skin disease	Patients with comorbidities
Chronicity of the skin disease	Patients whose diseases remit and recur
Skin disease impact on the patient’s quality of life	Patients with seemingly poor quality of life
Nature of the patient’s work	Patients with tight work schedules
Mobility of patient	Elderly patients or patients who live far from the hospital
Presence of a carer	Mainly in children or vulnerable patients
Patient-carer or patient-parent relationship	Mainly in children or vulnerable patients
Quality of the patient-GP relationship	Patients insisting on follow-up or unwilling to seek care from their GP
Local or national guidelines	When faced with clinical uncertainty
Appropriate management support from other care	Patients with comorbidities

providers such as oncologists, psychologists	
Healthcare service financial resources	Limited healthcare budget
Local discharge policy	Relevant clinic discharge policy
Referral of patient to nurse led clinics in secondary care	When patients need educational or treatment support

Table 4. Methods to train dermatology trainees how to discharge outpatients

Training Tips	Consultant quotations
Learning through modelling a senior consultant	<i>“Learning to discharge is by example rather than some written guideline.”</i>
Supervised “hands on” experience	<i>“You have to teach them how to discharge patients: as I have said before, whether they are safe to be discharged, are they stabilized, and is the GP able to monitor them, I think you’ve got to make a decision to discharge patients!”</i>
Review by consultant of every patient that is discharged by a trainee	<i>“I checked the [trainee’s] letters back to GPs, explaining how to give them a definite treatment plan, and suggesting what to do in this situation.”</i>
Learn to access and use relevant ‘Guidelines’	<i>“I guess if we were perhaps more fully supported with specific guidelines for specific conditions, in terms of exact follow up guidance, we felt supported by that.”</i>
Encourage self-confidence and acting on the basis of clinical judgement	<i>“If I really thought clinically that the patient did not really need to be coming back for a clinic appointment, I would make a real effort to discharge them. I would try to explain to them clearly why they didn’t need to come.”</i>
Collaborate and communicate well as a team	<i>“I guess this reflects on us as the supervising clinic, so good communication between juniors and seniors about which patients needs to be followed-up.”</i>

Table 5. Proposed educational objectives concerning the outpatient discharge decision.

Educational Objectives
To adopt a “shared decision-making” approach rather than a paternalistic approach
To self-recognise and manage poor verbal and non-verbal communication with patients
To recognise and manage ineffective collaboration with colleagues in primary and secondary care
To demonstrate confidence when making discharge decisions
To use the “Traffic-light” design discharge information checklist during training and routinely where appropriate
To evaluate strategies for appropriate discharge decisions
To recognise biases related to taking discharge decisions, such as overconfidence and anchoring biases
To manage consultation times to ensure sufficient time available for the discharge process
To reflect on the appropriateness of clinical decisions taken either to discharge or follow-up outpatients.

Legend to Figure:

Figure 1. Key aspects of an appropriate discharge