

**ORIGINAL RESEARCH:
EMPIRICAL RESEARCH - QUANTITATIVE**

Patient safety culture among European cancer nurses—An exploratory, cross-sectional survey comparing data from Estonia, Germany, Netherlands, and United Kingdom

Lena Sharp PhD, RN, Head of Department^{1,2} | Kristi Rannus PhD, RN, Head Nurse^{3,4} |
Anna Olofsson MSc, Statistician¹ | Daniel Kelly PhD, RN, Professor⁵ |
Wendy H. Oldenmenger PhD, RN, Coordinator oncology nursing research⁶ |
on behalf of the EONS RECaN group

¹Regional Cancer Centre, Stockholm-Gotland, Stockholm, Sweden

²Division of Innovative Care Research, Department of Learning Informatics, Management and Ethics, Karolinska Institutet, Stockholm, Sweden

³North Estonia Medical Centre, Tallinn, Estonia

⁴Tallinn Health Care College, Tallinn, Estonia

⁵School of Healthcare Sciences, Cardiff University, Wales, UK

⁶Department of Medical Oncology, Erasmus MC Cancer Institute, Rotterdam, The Netherlands

Correspondence

Wendy H. Oldenmenger, Department of Medical Oncology, Erasmus MC Cancer Institute, Rotterdam, The Netherlands.
Email: w.h.oldenmenger@erasmusmc.nl

Funding information

This project is jointly funded by the European Oncology Nursing Society (EONS) and the European Cancer Organisation (ECCO). In addition, the authors wanted to thank the Regional Cancer Centre Stockholm-Gotland for their additional funding for this phase 2 project.

[Correction added on 5 June 2020, after first online publication: Peer review history statement has been added.]

Abstract

Aim: To explore the differences in perceived patient safety culture in cancer nurses working in Estonia, Germany, the Netherlands, and the United Kingdom.

Design: An exploratory cross-sectional survey.

Methods: In 2018, 393 cancer nurses completed the 12 dimensions of the Hospital Survey on Patient Safety Culture.

Results: The mean score for the overall patient safety grade was 61.3. The highest rated dimension was “teamwork within units” while “staffing” was the lowest in all four countries. Nurses in the Netherlands and in the United Kingdom, scored higher on “communication openness”, the “frequency of events reported”, and “non-punitive response to errors”, than nurses from Estonia or Germany. We found statistically significant differences between the countries for the association between five of the 12 dimensions with the overall patient safety grade: overall perception of patient safety, communication openness, staffing, handoffs and transitions and non-punitive response to errors.

Conclusion: Patient safety culture, as reported by cancer nurses, varies between European countries and contextual factors, such as recognition of the nursing role and education have an impact on it. Cancer nurses' role in promoting patient safety is a key concern and requires better recognition on a European and global level.

Impact: Cancer Nursing Societies in any country can use these data as an indication on how to improve patient care in their country. Recognition of cancer nursing as a distinct speciality in nursing will help to improve patient safety.

KEYWORDS

cancer nurses, hospital survey on patient safety, nursing, patient safety culture

The peer review history for this article is available at <https://publons.com/publon/10.1111/JAN.14177>

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2019 The Authors. *Journal of Advanced Nursing* published by John Wiley & Sons Ltd.

1 | INTRODUCTION

Patient safety is defined as the prevention of errors and adverse effects to patients associated with health care (World Health Organisation, 2018). The World Health Organization (WHO) points out that health care has become more effective during recent decades, but also more complex and that these complexities may challenge efforts to improve patient safety as risks may increase.

Recent positive developments in cancer care with improved treatment outcomes also carry with them new patient safety risks. Potent drugs with small therapeutic margins, complex treatment regimens with severe symptom burdens and issues around adherence to treatment are examples of some of these risks (Weingart, Zhang, Sweeney, & Hassett, 2018). Patients with cancer are vulnerable to errors and mistakes can have catastrophic consequences (Weingart et al., 2018).

Research shows that errors occur during all phases of the administration process of cytotoxic drugs (Fyhr & Akselsson, 2012; Keers, Williams, Cooke, & Ashcroft, 2013; Kullberg, Larsen, & Sharp, 2013; Weingart et al., 2018). Cancer nurses have a central role in this process and they are often the last point of contact during this complex process prior to the drugs reaching the patient (Schwappach & Gehring, 2014). Nurses' actions and risk assessment skills are of great importance. Safe procedures and the correct use of devices are crucial steps in safety promotion (Kullberg et al., 2013; Mattsson et al., 2015), as is the courage to speak up and question when adverse events do occur in practice, including risks and near misses (Schwappach & Gehring, 2014). This requires a safety culture where adverse events can be reported without staff being blamed and if mistakes do occur, lessons are learned. Patient safety culture has been defined as the overall behaviour of individuals and organizations, based on a common set of beliefs and values that are aimed at reducing the opportunities for patient harm (Singer & Vogus, 2013). The terms "patient safety culture" and "patient safety climate" are sometimes used interchangeably. Safety climate has been described as a snapshot of the underlying safety culture (Danielsson, Nilsen, Rutberg, & Arestedt, 2017). The patient safety culture is an important measure in assessing the quality of health care and generally is measured by surveys (Danielsson et al., 2017; Mascherek & Schwappach, 2017). Research has shown that a high patient safety culture is associated with fewer readmission events (Hansen, Williams, & Singer, 2011), fewer medication errors and a reduction in urinary tract infections (Hofmann & Mark, 2006). Better perceptions of overall patient safety and a higher patient safety grade are also associated with nurses with higher levels of motivation (Toode, Routasalo, Helminen, & Suominen, 2015) and more satisfied patients and nurses (Hofmann & Mark, 2006).

1.1 | Background

Nurses are considered the most trusted profession in many countries (Brenan, 2018; Stephenson, 2018) and have a central role for people affected by cancer because they represent the largest

group of healthcare professionals in the cancer workforce. A recent systematic review (Charalambous et al., 2018) concluded that the contribution of cancer nursing to interventions to benefit patients and to cancer research more generally is significant but is not always recognized. The recognition of cancer nursing as a specialty across Europe is highly variable at present. The RECaN project (Recognising European Cancer Nursing) (Campbell et al., 2017; Charalambous et al., 2018; Kelly & Charalambous, 2017) has been initiated and conducted by the European Oncology Nursing Society (EONS) and supported by the European Cancer Organisation (ECCO). The overall goal is to increase recognition of the value and contribution of cancer nursing across Europe. This exploratory study is a part of the second phase of the RECaN project, where the patient safety culture in cancer nursing is compared across four European countries.

2 | THE STUDY

2.1 | Aim

The aim of this study was to explore the differences in the perceived patient safety culture in cancer nurses working in four European countries using the Hospital Survey on Patient Safety Culture (HSPSC). More specifically, the objectives were to compare perceptions and aspects of patient safety cultures that were important as described by cancer nurses in Estonia, Germany, the Netherlands, and the United Kingdom. Besides this goal, we also sought to identify those factors most significantly associated with the highest rates of overall patient safety assessments.

2.2 | Design and ethical considerations

We conducted an exploratory cross-sectional study to investigate workplace patient safety culture among cancer nurses in four European countries, i.e., Estonia, Germany, Netherlands, and the United Kingdom. The survey was conducted in all four countries during 2017. Research Ethics Committee approval for this study was obtained from the Ethical Review Committee at Cardiff University in the United Kingdom, with country specific approval from the Ethical Review Board of the Erasmus University Medical Center in the Netherlands and the Research Ethics Committee of the University of Tartu in Estonia. In Germany, additional approval was not necessary.

2.3 | Participants

Eligible participants were cancer nurses from the four countries involved. The data collection was conducted anonymously on a volunteer basis during annual conferences of the National Cancer Nursing Societies in each country, between May - November 2017. During the conferences, all cancer nurses had the opportunity to complete the questionnaire following an invitation from one of the researchers. This was voluntary and the nurses themselves decided whether they wished to participate. A specific database was developed for

the purpose of this study and data from all four countries were stored using anonymized codes.

2.4 | Data collection

The HSPSC assesses staff perceptions of the patient safety culture, including different aspects of safety, medical errors, and incident reporting. The survey is available in all four languages of the participating countries (Hammer et al., 2011; Smits et al., 2012; Smits, Wagner, Spreeuwenberg, van der Wal, & Groenewegen, 2009; Toode et al., 2015; Waterson, Griffiths, Stride, Murphy, & Hignett, 2010). The Agency for Healthcare Research and Quality gave permission to use the HSPSC in all four languages.

The HSPSC consists of some background variables, e.g. professional experience (years), work time (hours per week), primary working area and whether the nurses participating have direct patient contact. However, we could not collect these background data from Germany, since the German version of HSPSC did not include these variables.

The HSPSC includes 42 items covering 12 dimensions of the patient safety culture, with three or four items per dimension (File S1). All items are based on a five-point Likert-type scale, from "strongly disagree" to "strongly agree", or from "never" to "always". When necessary, prior to statistical analysis, negatively worded items were reverse coded so that a higher score always represented a positive response. The HSPSC also includes two single-items that provides an "overall patient safety grade", with a five-point Likert-type scale response (from "Failing" to "Excellent") and a "number of events reported" (Sorra et al., 2018). The overall patient safety grade item was used as an outcome variable in this study (Sorra & Dyer, 2010).

2.5 | Validity and reliability

The HSPSC has been published and showed acceptable psychometric properties, as factor analyses confirmed the validity of the HSPSC subscales and the questionnaire showed acceptable levels of reliability across the involved countries. Different studies showed a Cronbach's alpha $>.70$, which is acceptable, except for staffing (around 0.60 in most studies). All items of the HSPSC correlate significantly with the safety score (Blegen, Gearhart, O'Brien, Sehgal, & Alldredge, 2009; Lee, Phan, Dorman, Weaver, & Pronovost, 2016; Smits, Christiaans-Dingelhoff, Wagner, Wal, & Groenewegen, 2008).

2.6 | Data analysis

Prior to the statistical analysis, the scores of negatively worded items were reversed to ensure that higher scores always reflected a more positive assessment of patient safety culture. The dimension scores were then analysed using two different methods. (a) Percentages of positive responses, defined as values "agree" to "strongly agree" and "most of the time" to "always" (Sorra et al., 2018). The single item "overall patient safety grade" was dichotomized into high ("excellent" and "very good") and low ("acceptable", "fair", and "failing"). The single item "number of events reported" was dichotomized into no

events reported and one or more events reported. For each item, we calculated the percentages of respondents who answered positively. Then unweighted averages of those percentages were computed for each dimension, resulting in dimension scores ranging from 0–100; (b) The Likert-type scale was linearly transformed to a 100-point scale (Scaled score = [(Raw score – min response score)/range of possible response category scores]*100), with the lowest possible value corresponding to 0 and the highest possible value corresponding to 100 (Danielsson et al., 2017; Fayers et al., 2001).

Data were analysed using Statistical Program R. Demographic characteristics and the scores of the patient safety culture dimensions were summarized using descriptive statistics. Overall differences in the 12 dimensions between the four countries were tested using the Kruskal–Wallis test. If $p < .05$ was achieved the test was considered significant and a pairwise Wilcoxon test was performed to analyse any group difference. A multiple logistic regression analysis was performed to determine the association between the 12 dimensions and the four countries involved (explanatory variables), with countries \times dimension as an interaction term and the overall patient safety grade (outcome variable). When there was an overall difference ($p < .05$), a pairwise test was performed with a Bonferroni correction ($p < .01$ was considered significant).

3 | RESULTS

The sample in this study consisted of 393 European cancer nurses. Most of these cancer nurses worked in Germany ($N = 160$ [41%]), followed by the United Kingdom ($N = 94$ [24%]), the Netherlands ($N = 74$ [19%]) and Estonia ($N = 64$ [16%]). Most respondents worked in a Department of Medical Oncology ($N = 171$ [77%]), or Cancer Surgery ($N = 22$ [10%]). One third of respondents had professional experience of ≤ 5 years, one-third between 6–15 years and a final third had worked as a nurse for more than 15 years. Furthermore, most participants had worked <5 years in the current area (56%), had direct patient contact and worked less than 40 hr a week (60%, Table 1). Cronbach's alpha of each dimension was acceptable, except the dimension of "staffing" scored lower (0.53–0.66) in all four countries.

3.1 | Patient safety culture dimensions

The mean score for "overall patient safety grade" was 61.3 for the total sample, but this score varied statistically significantly between the countries, as tested with the Kruskal–Wallis test ($p < .0001$). Cancer nurses from Germany had the lowest score (mean 55.5), while cancer nurses from the United Kingdom scored the highest patient safety grade (mean 72.0, Table 2).

Overall, the highest rated patient safety culture dimensions were "teamwork within units" (mean 69.4), "organizational learning" (mean 64.8) and "feedback and communication about errors" (mean 64.5). The lowest rated dimensions were "handoffs and transitions" (mean 47.9) and "staffing" (mean 46.6, Table 2). Various dimensions of patient safety culture were rated differently in statistically significant ways between the four countries. Cancer nurses from the United

TABLE 1 Respondent characteristics

	All participants (N = 393)	Estonia (N = 64)	Germany (N = 160)	Netherlands (N = 74)	United Kingdom (N = 95)	p-value
Primary working area, n (%)						
Medical oncology	171 (77)	40 (63)	–	60 (87)	71 (81)	.0018
Surgery	22 (10)	11 (17)		7 (10)	4 (5)	
Others	28 (13)	13 (20)		2 (3)	13 (15)	
Professional experience, n (%), years						
≤5	83 (37)	22 (34)	–	25 (35)	36 (40)	.089
6–15	76 (34)	23 (36)		31 (44)	22 (24)	
>15	67 (30)	19 (27)		15 (21)	33 (36)	
Years in work area, n (%), years						
≤5	126 (56)	36 (56)	–	33 (46)	57 (63)	.111
6–15	65 (29)	21 (33)		26 (37)	18 (20)	
≥16	35 (15)	7 (11)		12 (17)	16 (18)	
Years in this hospital, n (%), years						
≤5	82 (36)	24 (38)	–	22 (31)	36 (40)	.019
6–15	71 (31)	27 (42)		17 (24)	27 (30)	
≥16	74 (33)	13 (20)		33 (46)	28 (31)	
Weekly work time, n (%), hours						
≤39	136 (60)	10 (16)	–	66 (93)	60 (66)	.002
>39	90 (40)	54 (84)		5 (7)	31 (34)	
Direct contact with patients, n(%)						
Yes	208 (93)	59 (92)	–	66 (97)	83 (91)	<.0001
No	15 (7)	5 (8)		2 (3)	8 (9)	

Note: p-value of the background characteristics were calculated with chi square test.

Kingdom scored higher than cancer nurses from the three other countries in most dimensions. This was especially evident on the dimensions “teamwork within units”, “supervisor/manager expectations and actions promoting patient safety”, “management support for patient safety” and “frequency of events reported” (Table 1). In all four countries, the dimension of staffing scored very low. This was mainly explained by the items that indicated that in all countries there were not enough cancer nurses available to handle the workload and that they were working in a “crisis mode” most of the time (File S1).

On the other hand, when looking at the percentages of positive answers overall, cancer nurses from the Netherlands scored the highest on “events reported”, “communication openness” and “non-punitive response to errors”. Whereas cancer nurses from the United Kingdom offered the most positive perceptions of the single item question “overall patient safety grade”. Cancer nurses from both the United Kingdom and the Netherlands scored higher on the dimension “frequency of events reported” than cancer nurses from Estonia or Germany (Figure 1).

3.2 | Factors associated with overall patient safety grade

Multiple logistic regression analysis showed a statistically significant difference between the four countries for associations between five

of 12 dimensions of patient safety culture with the overall patient safety grade: overall perception of patient safety, communication openness, staffing, handoffs and transitions and non-punitive response to errors (Table 3). A higher level of these five dimensions of patient safety culture implies increased probability for a higher overall patient safety grade. We found a statistically significant difference for the “overall perceptions of patient safety” between the four countries. This was mainly explained by the difference between Estonia and Germany (odds ratio [OR] 0.92, 95% confidence interval [CI] 0.865–0.969, pairwise Wilcoxon test: $p < .002$), meaning that for every increase in the “overall perceptions of patient safety” the change of having a higher “patient safety grade” was lower for Estonia compared with Germany. The statistically significant difference for “communication openness” was also explained by the differences between Estonia and Germany (OR [95% CI] = 0.93 [0.89–0.97], pairwise Wilcoxon test: $p < .001$). The statistically significant difference for “staffing” was explained by two comparisons, both between Estonia and Germany (OR [95% CI] = 0.92 [0.88–0.96], pairwise Wilcoxon test: $p < .001$) and the United Kingdom and Germany (OR [95% CI] = 0.95 [0.91–0.99], pairwise Wilcoxon test: $p < .001$). While the difference for “handoffs and transitions” was mainly explained by the difference between the Netherlands and Germany (OR [95% CI] = 0.93 [0.89–0.98], pairwise Wilcoxon

TABLE 2 Distribution of the dimensions of the HSPSC

	All participants (N = 393)	Estonia (N = 64)	Germany (N = 160)	Netherlands (N = 74)	United Kingdom (N = 95)	p-value
Patient safety culture dimensions, mean (SD)						
Teamwork within units	69.4 (17.0)	69.2 (19.1)	65.1 (16.2)	72.0 (15.3)	74.8 (16.0)	<.0001
Supervisor/manager expectations and actions promoting safety	63.0 (20.1)	67.8 (17.2)	59.2 (22.2)	56.9 (15.8)	72.0 (17.2)	<.0001
Organizational learning	64.8 (16.0)	66.7 (13.9)	62.3 (17.2)	62.6 (13.3)	69.5 (16.2)	.0004
Management support for patient safety	54.5 (19.8)	59.0 (17.6)	48.5 (22.1)	54.8 (6.1)	61.7 (20.7)	<.0001
Overall perception of patient safety	58.2 (17.6)	61.6 (15.5)	54.6 (17.4)	60.3 (17.5)	60.7 (18.4)	.003
Feedback and communication about errors	64.5 (19.8)	63.0 (20.2)	62.1 (20.8)	67.0 (14.9)	67.6 (18.4)	.156
Communication openness	63.1 (17.6)	65.6 (17.5)	60.5 (18.6)	60.1 (14.9)	67.5 (17.2)	.006
Frequency of events reported	56.7 (23.6)	44.6 (23.1)	51.1 (22.9)	63.0 (18.9)	69.5 (21.0)	<.0001
Teamwork across units	54.6 (16.1)	58.0 (14.8)	55.1 (14.7)	52.1 (18.9)	53.2 (16.6)	.223
Staffing	46.6 (18.6)	48.0 (19.1)	43.8 (18.9)	47.8 (15.1)	49.6 (19.1)	.178
Handoffs and transitions	47.9 (16.1)	46.4 (14.1)	48.1 (15.4)	45.2 (13.4)	51.0 (19.7)	.045
Non-punitive response to errors	57.7 (19.7)	56.6 (18.9)	60.1 (20.0)	53.2 (14.6)	57.7 (22.7)	.021
Overall patient safety grade, mean (SD)	61.3 (18.7)	61.3 (16.6)	55.5 (18.6)	60.4 (15.6)	72.0 (17.8)	<.0001

Note: p-value patient safety culture was calculated with Kruskal-Wallis test.

test: $p < .006$). In the multiple logistic regression analysis, we found a $p = .024$ for “non-punitive response to errors”. However, all pairwise comparisons showed a $p > .01$, due to the Bonferroni correction, meaning there is no significant difference between any of the countries for this variable.

4 | DISCUSSION

To the best of our knowledge, this is the first study comparing patient safety culture perceptions between cancer nurses in Estonia, Germany, the Netherlands and the United Kingdom. We demonstrated a statistically significant difference in the “overall patient safety grade” between the four countries. This was mainly explained by a statistically significant difference in the following dimensions of safety culture as measured by the HSPSC; overall perception of patient safety, communication openness, staffing, handoffs and transitions and non-punitive response to errors.

One of the most important factors associated with a positive patient safety culture is openness in communication. This is not just about communication itself, but also related to nurses feeling confident to speak-up and report unsafe events, without fear of negative consequences. Consequently, as team members nurses have the

opportunity to learn from both their own and other’s mistakes, fostering a more open climate of organizational learning. In the current study, the nurses in the Netherlands and in the UK, scored higher on “communication openness”, the “frequency of events reported” and “non-punitive response to errors” (Figure 1 and File S1). This is contrary with existing literature, where respondents in other studies scored remarkably lower on these items (Alswat et al., 2017; Blegen et al., 2009; Famolaro et al., 2016; Mardon, Khanna, Sorra, Dyer, & Famolaro, 2010). Our results may reflect the differences between the four countries in our study, especially in terms of contextual factors such as; recognition, autonomy, career opportunities, and educational preparation. Cancer nursing in both the Netherlands and the United Kingdom is recognized professionally and is well-established as a distinct specialty in nursing. Both countries also have career possibilities in clinical cancer nursing, e.g. advanced nursing roles. However, in the other two countries this is not currently the case.

Patient safety is one of the most important factors in quality of care and is inseparable from the safety culture (Charalambous & Kelly, 2018; Ulrich & Kear, 2014). In the current study, one of the statistically significant factors related to patient safety culture was staffing. The percentage of positive responses, however, are lower for all staffing items, in all four countries (File S1; Figure 1). Earlier




FIGURE 1 Percentage of the positive scores of the patient safety culture dimensions and single-items per country

Dimensions	p^a
Teamwork within units	.087
Supervisor/manager expectations and actions promoting safety	.097
Organizational learning	.301
Management support for patient safety	.150
Overall perception of patient safety	.003
Feedback and communication about errors	.073
Communication openness	.007
Frequency of events reported	.138
Teamwork across units	.074
Staffing	<.0001
Handoffs and transitions	.022
Non-punitive response to errors	.024

^aMultiple logistic regression analysis.

TABLE 3 The association between the dimensions of the HSPSC and countries (explanatory variables) and overall patient safety grade

studies from the Arabic countries reported similarly low numbers (Alswat et al., 2017; El-Jardali, Jaafar, Dimassi, Jamal, & Hamdan, 2010; Hamdan & Saleem, 2013), in contrast with studies from the USA and Sweden where respondents scored considerably higher regarding staffing (Blegen et al., 2009; Danielsson et al., 2017; Famolaro et al., 2016; Mardon et al., 2010). In our study, nurses from the Netherlands and the United Kingdom may be questioning whether there are enough nurses to guarantee patient safety, while in the other countries, the question is not only one of numbers, but also of the availability of suitably qualified specialist cancer nurses to promote a safety culture.

Earlier studies, such as the RN4CAST project, showed that nurses' workload and level of education were directly linked with patient outcomes and, ultimately, with patient mortality (Aiken et al., 2014, 2017). Cancer nursing is developing to meet the rising demands of increased cancer incidence, prevalence and newer and more complex treatment options. The need for expertise in specialized and advanced cancer nursing is therefore also increasing (Charalambous et al., 2018). Some European countries have already implemented and seen the benefits from advanced cancer nursing roles to start meeting these rising needs (Cowman et al., 2010), while other do not yet recognize advanced nursing roles or

even offer postgraduation education. To help countries to establish specialized education in cancer nursing, EONS published a Cancer Nursing Education Framework (2018), which describes the competences involved in cancer nursing and how these can be addressed in education programs globally.

4.1 | Limitations

Although we used a validated questionnaire to measure patient safety culture, it is not always possible to compare these data to other studies. Most other studies include a variety of respondents such as managers, physicians, technicians, and sometimes but not always, nurses (Danielsson et al., 2017; Sorra et al., 2018). More importantly, such data are often reported in different ways. In some studies, the means of the 12 HSPSC dimensions are described as the mean of the Likert-type scale (Burlison et al., 2016; Hammer et al., 2011; Smits et al., 2012), or a transformation of the Likert-type scale into a 0–100 scale (Table 1; (Danielsson et al., 2017)). The disadvantage of using a mean is that it tends to shift towards the middle. Therefore, it does not always correctly reflect the range of opinions of all the respondents. In other studies, the percentage of positive answers for the 12 dimensions were reported (Figure 1; Alswat et al., 2017; Danielsson et al., 2017; Sorra et al., 2018), which may give a clearer reflection of the given responses. Our study was also carried out with a self-selecting sample and may reflect the views of those motivated to attend professional conferences. Furthermore, we could not report on response rates. Wider samples and other recruitment strategies should be considered in future safety culture research studies involving cancer nurses.

This study presents an analysis of cancer nurses' opinions about the patient safety cultures in four European countries. Therefore, the results can only be seen as an indication about patient safety culture in these settings. Nevertheless, our data are derived from the responses of cancer nurses with a range of professional expertise and considerable years of experience of working in their current workplaces. As much as 93% of our respondents worked in direct contact with patients, representing what we suggest is a "true" range of clinical cancer nurses. Based on this study, we suggest that organizations need to ensure that cancer nurses are recognized as key members of multi-professional care teams. This will require the further development of the status and recognition of cancer nursing in all countries to enhance the patient safety culture in cancer care.

5 | CONCLUSION

This study showed that cancer nurses scored patient safety differently between four European countries. This is mainly explained by the dimensions of overall perception of: patient safety, communication openness, staffing, handoffs and transitions and non-punitive response to errors.

As nurses are the largest group of professionals working directly with cancer patients the growing rates of cancer incidence

and prevalence will increase the need for specialized cancer nursing roles in all countries. Nurses work closely with patients across the entire cancer trajectory and are therefore the best suited professionals to assess and promote the patient safety culture, however, all health professionals also have a responsibility in this regard.

As well as better awareness of patient safety generally there are also other issues now to be considered in practice contexts. These include having adequate numbers of Registered Nurses to meet the safety needs of patients with cancer. More research is now needed to examine cancer patient acuity in relation to nurse staffing levels and other factors such as the experience level and qualifications of the nurses available. Evidence already exists from international studies of the link between nurse staffing and nurse qualifications with patient mortality in acute medical and surgical units (Aiken et al., 2014, 2017).

Research effort should now be extended to oncology settings with agreed nurse-specific outcomes being identified that reflect the needs of cancer patients such as infection, nausea, and vomiting or pain (Aiken et al., 2017; Oldenmenger et al., 2018). By doing so the safety culture of cancer settings will be enhanced by improving the visibility of the contribution made by cancer nurses and ensuring that they feel confident about speaking up when safety concerns arise.

ACKNOWLEDGEMENTS

The authors thank the four national cancer nursing societies: Estonia Oncology Nursing Society, KOK, UKON, V&VN Oncologie, and the other members of the RECaN phase 2 group for their contributions: Patrick Crombez, Patrick Jahn, Gabi Knoetgen, Annette Lanskshear, Theresa Wiseman, and Gaby Wydenkeller.

CONFLICTS OF INTEREST

No conflict of interest has been declared by the authors.

AUTHOR CONTRIBUTIONS

All authors made substantial contributions to conception and design, or acquisition of data, or analysis and interpretation of data; involved in drafting the manuscript or revising it critically for important intellectual content; given final approval of the version to be published. Each author should have participated sufficiently in the work to take public responsibility for appropriate portions of the content; agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

ORCID

Lena Sharp  <https://orcid.org/0000-0002-0818-6992>

Daniel Kelly  <https://orcid.org/0000-0002-1847-0655>

Wendy H. Oldenmenger  <https://orcid.org/0000-0001-6855-6505>

REFERENCES

- Aiken, L. H., Sloane, D. M., Bruyneel, L., Van den Heede, K., Griffiths, P., Busse, R., ... RN4CAST consortium. (2014). Nurse staffing and education and hospital mortality in nine European countries: A retrospective observational study. *Lancet*, *383*(9931), 1824–1830.
- Aiken, L. H., Sloane, D., Griffiths, P., Rafferty, A. M., Bruyneel, L., McHugh, M. ... RN4CAST consortium. (2017). Nursing skill mix in European hospitals: Cross-sectional study of the association with mortality, patient ratings and quality of care. *BMJ Quality Safety*, *26*(7), 559–568. <https://doi.org/10.1136/bmjqs-2016-005567>
- Alswat, K., Abdalla, R. A. M., Titi, M. A., Bakash, M., Mehmood, F., Zubairi, B., ... El-Jardali, F. (2017). Improving patient safety culture in Saudi Arabia (2012–2015): Trending, improvement and benchmarking. *BMC Health Services Research*, *17*(1), 516. <https://doi.org/10.1186/s12913-017-2461-3>
- Blegen, M. A., Gearhart, S., O'Brien, R., Sehgal, N. L., & Alldredge, B. K. (2009). AHRQ's hospital survey on patient safety culture: Psychometric analyses. *Journal of Patient Safety*, *5*(3), 139–144. <https://doi.org/10.1097/PTS.0b013e3181b53f6e>
- Brenan, M. (2018). *Nurses again outpace other professionals for honesty, ethics*. Retrieved from <https://news.gallup.com/poll/245597/nurses-again-outpace-professions-honesty-ethics.aspx>
- Burlison, J. D., Quillivan, R. R., Kath, L. M., Zhou, Y., Courtney, S. C., Cheng, C., & Hoffman, J. M. (2016). A multilevel analysis of U.S. hospital patient safety culture relationships with perceptions of voluntary event reporting. *Journal of Patient Safety* <https://doi.org/10.1097/PTS.0000000000000336>
- Campbell, P., Torrens, C., Kelly, D., Charalambous, A., Domenech-Climent, N., Nohavova, I., ... Wells, M. (2017). Recognizing European cancer nursing: Protocol for a systematic review and meta-analysis of the evidence of effectiveness and value of cancer nursing. *Journal of Advanced Nursing*, *73*(12), 3144–3153. <https://doi.org/10.1111/jan.13392>
- Charalambous, A., & Kelly, D. (2018). Promoting a safety culture through effective nursing leadership in cancer care. *European Journal of Oncology Nursing*, *36*, vi–vii. <https://doi.org/10.1016/j.ejon.2018.10.002>
- Charalambous, A., Wells, M., Campbell, P., Torrens, C., Östlund, U., Oldenmenger, W., ... Kelly, D. (2018). A scoping review of trials of interventions led or delivered by cancer nurses. *International Journal of Nursing Studies*, *86*, 36–43. <https://doi.org/10.1016/j.ijnurstu.2018.05.014>
- Cowman, S., Gethin, G., O'Neill, M., Lawrence, S., Kinsella, K., Lavelle, A., & Roberts, A. (2010). *Evaluation of the role of the clinical nurse specialist in cancer care*. Dublin: Centre for Nursing and Midwifery Research, Royal College of Surgeons in Ireland. Retrieved from http://www.rcsi.ie/files/facultyofnursingmidwifery/docs/20110518024459_Final%20Report%20CNS%20Study%20July%2020.pdf
- Danielsson, M., Nilsen, P., Rutberg, H., & Arestedt, K. (2017). A national study of patient safety culture in hospitals in Sweden. *Journal of Patient Safety*, <https://doi.org/10.1097/PTS.0000000000000369>
- El-Jardali, F., Jaafar, M., Dimassi, H., Jamal, D., & Hamdan, R. (2010). The current state of patient safety culture in Lebanese hospitals: A study at baseline. *International Journal for Quality in Health Care*, *22*(5), 386–395. <https://doi.org/10.1093/intqhc/mzq047>
- EONS Cancer Nursing Education Framework. (2018). Retrieved from Brussels: <http://www.cancernurse.eu/documents/EONSCancerNursingFramework2018.pdf>
- Famolaro, M. P. S., Dyer, N., Burns, W., Flashner, E., Liu, H., & Sorra, J. (2016). Hospital survey on patient safety culture 2016 user comparative database report (No. 16–0021-EF). Retrieved from https://www.ahrq.gov/sites/default/files/wysiwyg/professionals/quality-patient-safety/patientsafetyculture/hospital/2016/2016_hospitalops_report_pt1.pdf
- Fayers, P. M., Aaronson, N. K., Bjordal, K., Grounvoid, M., Curran, D., & Bottomley, A. (2001). *The EORTC QLQ-C30 scoring manual* (3rd ed.). Brussels: European Organisation for Research and Treatment of Cancer.
- Fyhr, A., & Akselsson, R. (2012). Characteristics of medication errors with parenteral cytotoxic drugs. *European Journal of Cancer Care*, *21*(5), 606–613. <https://doi.org/10.1111/j.1365-2354.2012.01331.x>
- Hamdan, M., & Saleem, A. A. (2013). Assessment of patient safety culture in Palestinian public hospitals. *International Journal for Quality in Health Care*, *25*(2), 167–175. <https://doi.org/10.1093/intqhc/mzt007>
- Hammer, A., Ernstmann, N., Ommen, O., Wirtz, M., Manser, T., Pfeiffer, Y., & Pfaff, H. (2011). Psychometric properties of the Hospital Survey on Patient Safety Culture for hospital management (HSOPS_M). *BMC Health Services Research*, *11*, 165. <https://doi.org/10.1186/1472-6963-11-165>
- Hansen, L. O., Williams, M. V., & Singer, S. J. (2011). Perceptions of hospital safety climate and incidence of readmission. *Health Services Research*, *46*(2), 596–616. <https://doi.org/10.1111/j.1475-6773.2010.01204.x>
- Hofmann, D. A., & Mark, B. (2006). An investigation of the relationship between safety climate and medication errors as well as other nurse and patient outcomes. *Personnel Psychology*, *59*(4), 847–869. <https://doi.org/10.1111/j.1744-6570.2006.00056.x>
- Keers, R. N., Williams, S. D., Cooke, J., & Ashcroft, D. M. (2013). Causes of medication administration errors in hospitals: A systematic review of quantitative and qualitative evidence. *Drug Safety*, *36*(11), 1045–1067. <https://doi.org/10.1007/s40264-013-0090-2>
- Kelly, D., & Charalambous, A. (2017). Recognising the impact and future potential of cancer nursing: The RECaN project and beyond. *European Journal of Oncology Nursing*, *29*, A1–A2. <https://doi.org/10.1016/j.ejon.2017.06.004>
- Kullberg, A., Larsen, J., & Sharp, L. (2013). 'Why is there another person's name on my infusion bag?' Patient safety in chemotherapy care - a review of the literature. *European Journal of Oncology Nursing*, *17*(2), 228–235. <https://doi.org/10.1016/j.ejon.2012.07.005>
- Lee, S. H., Phan, P. H., Dorman, T., Weaver, S. J., & Pronovost, P. J. (2016). Handoffs, safety culture and practices: Evidence from the hospital survey on patient safety culture. *BMC Health Services Research*, *16*, 254. <https://doi.org/10.1186/s12913-016-1502-7>
- Mardon, R. E., Khanna, K., Sorra, J., Dyer, N., & Famolaro, T. (2010). Exploring relationships between hospital patient safety culture and adverse events. *Journal of Patient Safety*, *6*(4), 226–232. <https://doi.org/10.1097/PTS.0b013e3181fd1a00>
- Mascherek, A. C., & Schwappach, D. L. B. (2017). Patient safety climate profiles across time: Strength and level of safety climate associated with a quality improvement program in Switzerland-A cross-sectional survey study. *PLoS ONE*, *12*(7), e0181410. <https://doi.org/10.1371/journal.pone.0181410>
- Mattsson, T. O., Holm, B., Michelsen, H., Knudsen, J. L., Brixen, K., & Herrstedt, J. (2015). Non-intercepted dose errors in prescribing anti-neoplastic treatment: A prospective, comparative cohort study. *Annals of Oncology*, *26*(5), 981–986. <https://doi.org/10.1093/annonc/mdv032>
- Oldenmenger, W. H., Geerling, J. I., Mostovaya, I., Vissers, K. C. P., de Graeff, A., Reyners, A. K. L., & Van der Linden, Y. M. (2018). A systematic review of the effectiveness of patient-based educational interventions to improve cancer-related pain. *Cancer Treatment Reviews*, *63*, 96–103. <https://doi.org/10.1016/j.ctrv.2017.12.005>
- Schwappach, D. L., & Gehring, K. (2014). Trade-offs between voice and silence: A qualitative exploration of oncology staff's decisions to speak up about safety concerns. *BMC Health Services Research*, *14*, 303. <https://doi.org/10.1186/1472-6963-14-303>
- Singer, S. J., & Vogus, T. J. (2013). Reducing hospital errors: Interventions that build safety culture. *Annual Review Public Health*, *34*, 373–396. <https://doi.org/10.1146/annurev-publhealth-031912-114439>

- Smits, M., Christiaans-Dingelhoff, I., Wagner, C., Wal, G., & Groenewegen, P. P. (2008). The psychometric properties of the 'Hospital Survey on Patient Safety Culture' in Dutch hospitals. *BMC Health Services Research*, 8, 230. <https://doi.org/10.1186/1472-6963-8-230>
- Smits, M., Wagner, C., Spreeuwenberg, P., Timmermans, D. R., van der Wal, G., & Groenewegen, P. P. (2012). The role of patient safety culture in the causation of unintended events in hospitals. *Journal of Clinical Nursing*, 21(23–24), 3392–3401. <https://doi.org/10.1111/j.1365-2702.2012.04261.x>
- Smits, M., Wagner, C., Spreeuwenberg, P., van der Wal, G., & Groenewegen, P. P. (2009). Measuring patient safety culture: An assessment of the clustering of responses at unit level and hospital level. *Quality Safety Health Care*, 18(4), 292–296. <https://doi.org/10.1136/qshc.2007.025965>
- Sorra, J. S., & Dyer, N. (2010). Multilevel psychometric properties of the AHRQ hospital survey on patient safety culture. *BMC Health Services Research*, 10, 199. <https://doi.org/10.1186/1472-6963-10-199>
- Sorra, J., Gray, L., Streagle, S., Famolaro, T., Yount, N., & Behm, J. (2018). AHRQ hospital survey on patient safety culture: User's guide (AHRQ Publication No. 18-0036-EF (Replaces 04-0041, 15(16)-0049-EF)). Retrieved from <https://www.ahrq.gov/sites/default/files/wysiwyg/professionals/quality-patient-safety/patientsafetyculture/hospital/userguide/hospcult.pdf>
- Stephenson, J. (2018). *Nursing once again tops chart of most trusted professions*. Retrieved from <https://www.nursingtimes.net/news/reviews-and-reports/nursing-once-again-tops-chart-of-most-trusted-professions/7026833.article>
- Toode, K., Routasalo, P., Helminen, M., & Suominen, T. (2015). Hospital nurses' working conditions in relation to motivation and patient safety. *Nursing Management (Harrow)*, 21(10), 31–41. <https://doi.org/10.7748/nm.21.10.31.e1293>
- Ulrich, B., & Kear, T. (2014). Patient safety and patient safety culture: Foundations of excellent health care delivery. *Nephrology Nursing Journal*, 41(5), 447–456.
- Waterson, P., Griffiths, P., Stride, C., Murphy, J., & Hignett, S. (2010). Psychometric properties of the Hospital Survey on Patient Safety Culture: Findings from the UK. *Quality Safety Health Care*, 19(5), e2. <https://doi.org/10.1136/qshc.2008.031625>
- Weingart, S. N., Zhang, L., Sweeney, M., & Hassett, M. (2018). Chemotherapy medication errors. *Lancet Oncology*, 19(4), e191–e199. [https://doi.org/10.1016/S1470-2045\(18\)30094-9](https://doi.org/10.1016/S1470-2045(18)30094-9)
- World Health Organisation (2018). *Patient safety*. Retrieved from <http://www.euro.who.int/en/health-topics/Health-systems/patient-safety>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Sharp L, Rannus K, Olofsson A, Kelly D, Oldenmenger WH; on behalf of the EONS RECaN group. Patient safety culture among European cancer nurses—An exploratory, cross-sectional survey comparing data from Estonia, Germany, Netherlands, and United Kingdom. *J Adv Nurs*. 2019;75:3535–3543. <https://doi.org/10.1111/jan.14177>

The *Journal of Advanced Nursing (JAN)* is an international, peer-reviewed, scientific journal. *JAN* contributes to the advancement of evidence-based nursing, midwifery and health care by disseminating high quality research and scholarship of contemporary relevance and with potential to advance knowledge for practice, education, management or policy. *JAN* publishes research reviews, original research reports and methodological and theoretical papers.

For further information, please visit *JAN* on the Wiley Online Library website: www.wileyonlinelibrary.com/journal/jan

Reasons to publish your work in *JAN*:

- **High-impact forum:** the world's most cited nursing journal, with an Impact Factor of 1.998 – ranked 12/114 in the 2016 ISI Journal Citation Reports © (Nursing (Social Science)).
- **Most read nursing journal in the world:** over 3 million articles downloaded online per year and accessible in over 10,000 libraries worldwide (including over 3,500 in developing countries with free or low cost access).
- **Fast and easy online submission:** online submission at <http://mc.manuscriptcentral.com/jan>.
- **Positive publishing experience:** rapid double-blind peer review with constructive feedback.
- **Rapid online publication in five weeks:** average time from final manuscript arriving in production to online publication.
- **Online Open:** the option to pay to make your article freely and openly accessible to non-subscribers upon publication on Wiley Online Library, as well as the option to deposit the article in your own or your funding agency's preferred archive (e.g. PubMed).