

Supporting Information

Efficient Elimination of Chlorinated Organics on A Phosphoric Acid Modified CeO₂ Catalyst: A Hydrolytic Destruction Route

Xiaoxia Dai, Xinwei Wang, Samuel Pattison, Yunhao Lu, David J. Morgan, Stuart H. Taylor,

James H. Carter*, Graham J. Hutchings, Xiaole Weng*, Zhongbiao Wu

Number of Pages: 5

Number of Figures: 4

Number of Tables: 3




Figure S1. Optimized model of CeO₂ (110) with oxygen vacancy. The models were constructed by removing single oxygen atom from CeO₂(110) supercell to introduce oxygen vacancies. During geometry optimization, the atoms in the top two layers of CeO₂ slab were allowed to relax while atoms in the bottom two layers were fixed in their optimized bulk positions.




Figure S2. TEM images of CeO₂ and HP-CeO₂

Table S1. Physical properties of HP-CeO₂ and CeO₂

sample	p loading		surface area (m ² g ⁻¹)	lattice parameter (Å)	Ce ³⁺ /Ce _{total} ^a
	(mmol g ⁻¹)	(nm ⁻²)			
CeO₂	0	0	96	5.411	29%
HP-CeO₂	0.21	1.18	108	5.413	26%

Note: (a) caculated from XPS results




Figure S3. Stability test of CB oxidation over HP-CeO₂ at 230 °C. Reaction condition: GHSV at 10,000 mL/(g h), chlorobenzene at ca. 500 ppm, H₂O at 5000 ppm, N₂ flow rate at ca. 145 mL/min, O₂ flow rate at ca. 15 mL/min.




Figure S4. *in-situ* FTIR spectra of CB oxidation at 150 °C over CeO₂ catalyst in dry condition.

Table S2. Adsorption energies on various active sites

Species	Adsorption energy (eV)
O ₂ /O _{vac}	2.20
H ₂ O/O _{vac}	1.57
O ₂ +H ₂ O/O _{vac}	2.82
C ₆ H ₅ Cl/O _{vac}	0.39
C ₆ H ₅ Cl+O ₂ /O _{vac}	2.37

C ₆ H ₅ Cl+H ₂ O/O _{vac}	1.21
O ₂ /Ce	0.20
H ₂ O/Ce	0.62
O ₂ +H ₂ O/Ce	0.72
C ₆ H ₅ Cl/Ce	0.48
C ₆ H ₅ Cl+H ₂ O/Ce	1.81
O ₂ /P group	-1.59
H ₂ O/P group	0.63
O ₂ +H ₂ O/P group	-0.94
C ₆ H ₅ Cl/P group	0.50
C ₆ H ₅ Cl+O ₂ /P group	-0.56
C ₆ H ₅ Cl+H ₂ O/P group	2.77
HCl/P group	0.74
C ₆ H ₅ OH/P group	0.76


Table S3. Ingredients of 17 toxic dioxins in the off-gas of CeO₂ and HP-CeO₂ at 250 °C test with H₂O stream

Compound	Detection limit (pg)	TEF	CeO ₂		HP-Ce	
			Cout (pg)	I-TEQ (ng/m ³)	Cout (pg)	I-TEQ (ng/m ³)
2378TCDD	0.5627	1	<0.5627	ND	<0.5627	ND
12378PeCDD	0.6234	0.5	<0.6234	ND	<0.6234	ND
123478HxCDD	0.6574	0.1	<0.6574	ND	<0.6574	ND


123678HxCDD	0.625	0.1	1.4879	0.00155	<0.625	ND
123789HxCDD	0.6931	0.1	<0.6931	ND	<0.6931	ND
1234678HpCDD	0.5632	0.01	6.2856	0.000655	<0.5632	ND
OCDD	0.7633	0.001	4.2767	4.45E-05	<0.7633	ND
2378TCDF	0.4675	0.1	0.9775	0.001018	<0.4675	ND
12378PeCDF	0.642	0.05	1.2645	0.000659	<0.642	ND
23478PeCDF	0.6132	0.5	1.04855	0.005461	<0.6132	ND
123478HxCDF	0.6015	0.1	2.3201	0.002417	<0.6015	ND
123678HxCDF	0.6329	0.1	2.5632	0.00267	<0.6329	ND
123789HxCDF	0.6512	0.1	0.57475	0.000599	<0.6512	ND
234678HxCDF	0.6877	0.1	2.3671	0.002466	<0.6877	ND
1234678HpCDF	0.7022	0.01	5.3266	0.000555	<0.7022	ND
1234789HpCDF	0.6934	0.01	<0.6934	ND	<0.6934	ND
OCDF	0.7142	0.001	3.3276	3.47E-05	<0.7142	ND
Total				0.01813		ND

Note: (1) CDD/CDF: chlorinated dibenzo-*p*-dioxin/dibenzofuran, structural formula represents as Fig S10 ; (2) ND: not detectable; (3) I-TEQ = Cout × TEF; (4) TEF: Toxic Equivalent Factors, toxic factors relative to the most toxic congener, 2,3,7,8-tetrachlorodibenzo-dioxin; (5) I-TEQ: International Toxicity Equivalence Quotient, the weighted value of the concentrations of 17 PCDD and PCDF congeners with chlorine in the 2, 3, 7 and 8 positions on the dibenzo skeleton, weighted according their Toxic Equivalent Factors (TEF) relative to the most toxic congener, 2,3,7,8-tetrachlorodibenzo-dioxin.

Polychlorinated dibenzofurans



Polychlorinated dibenzo-*p*-dioxins



Structure diagrams of chlorinated dibenzo-*p*-dioxin/dibenzofuran