

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <http://orca.cf.ac.uk/72435/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Wanger, Tim, Dewitt, Sharon, Collins, Anne, Maitland, Norman J., Poghosyan, Zaruhi and Knauper, Vera 2015. Differential regulation of TROP2 release by PKC isoforms through vesicles and ADAM17. *Cellular Signalling* 27 (7) , pp. 1325-1335. 10.1016/j.cellsig.2015.03.017 filefile

Publishers page: <http://dx.doi.org/10.1016/j.cellsig.2015.03.017>
<<http://dx.doi.org/10.1016/j.cellsig.2015.03.017>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.






Fig. S1. TROP2 expression constructs and analysis of TROP2 levels using TROP2 antibody in non-reducing conditions.

(A) Increase in 11 kDa MP fragment in cell lysates in the presence or absence of 1 µM GW64 or GI23. Data were normalised to GAPDH loading control. N = 3 blots. (B) Analysis of AP-TROP2 levels in medium following ultracentrifugation using anti TROP2 antibody shows that AP-TROP2-ECD dimers are enriched in the soluble fraction. (C) Analysis of aMP fragment production in response to mPS treatment over time from the shedding data presented in Fig. 4A. by Western blotting using V5 antibody. (D) Schematic representation of TROP2 and AP-TROP2 expression constructs.




Fig. S2. Comparison of aMP-fragments in AP-TROP2, TROP2-V5 and PC3 cells following deglycosylation.

(A) AP-TROP2 cells were stimulated with mPS for 3 h prior to deglycosylation with PNGase F and analysed with V5-epitope antibody for the aMP fragment. Deglycosylation of the aMP fragment reduces its molecular mass to 32 kDa. (B) The same experiment described in A but performed with TROP2-V5 expressing cells shows induction of aMP fragment formation with mPS, showing an identical molecular weight after deglycosylation with PNGaseF. (C) LNCaP cell lysates contain fully and partially glycosylated TROP2 but lack the aMP fragment. (D) PC3 cell lysates are positive for aMP TROP2 fragment. (E) TROP2 is expressed in prostate cancer stem cells but not in benign hyperplasia (BPH sample). Samples in this blot represent glycosylated TROP2. Gleason score is indicated above and GAPDH loading control is shown below.