

Rescue of failed oocyte activation after ICSI in a mouse model of male factor infertility by recombinant phospholipase C ζ

Randa Sanusi¹, Yuansong Yu^{1,2}, Michail Nomikos¹, F. Anthony Lai¹, and Karl Swann^{1,*}

¹Institute of Molecular and Experimental Medicine, Cardiff University School of Medicine, Heath Park, Cardiff CF14 4XN, UK

²Present address: TopChoice Medical, TopChoice Building, 327 Tianmushan Road, Hangzhou City, Zhejiang Province, China

*Correspondence address. E-mail: swannk1@cf.ac.uk

Submitted on February 24, 2015; resubmitted on June 30, 2015; accepted on July 13, 2015

ABSTRACT: Artificial oocyte activation to overcome failed fertilization after intracytoplasmic sperm injection (ICSI) in human oocytes typically employs Ca²⁺ ionophores to produce a single cytosolic Ca²⁺ increase. In contrast, recombinant phospholipase C ζ (PLC ζ) causes Ca²⁺ oscillations indistinguishable from those occurring during fertilization, but remains untested for its efficacy in a scenario of ICSI fertilization failure. Here, we compare PLC ζ with other activation stimuli in a mouse model of failed oocyte activation after ICSI, in which heat-treated sperm are injected into mouse oocytes. We show that increasing periods of 56°C exposure of sperm produces a progressive loss of Ca²⁺ oscillations after ICSI. The decrease in Ca²⁺ oscillations produces a reduction in oocyte activation and embryo development to the blastocyst stage. We treated such oocytes that failed to activate after ICSI either with Ca²⁺ ionophore, or with Sr²⁺ media which causes Ca²⁺ oscillations, or we injected them with recombinant human PLC ζ . All these treatments rescued oocyte activation, although Sr²⁺ and PLC ζ gave the highest rates of development to blastocyst. When recombinant PLC ζ was given to oocytes previously injected with control sperm, they developed normally to the blastocyst stage at rates similar to that after control ICSI. The data suggest that recombinant human PLC ζ protein is an efficient means of rescuing oocyte activation after ICSI failure and that it can be effectively used even if the sperm already contains endogenous Ca²⁺ releasing activity.

Key words: PLC ζ / ICSI / fertilization / sperm / oocyte

Introduction

For couples attending *in vitro* fertilization (IVF) clinics, intracytoplasmic sperm injection (ICSI) is a widely used treatment option (Palermo *et al.*, 1992; Johnson *et al.*, 2013) that is remarkably successful, compared with conventional IVF treatments, for treating many cases of male factor infertility (Palermo *et al.*, 1992; Johnson *et al.*, 2013). However, cases of ICSI where all of the available oocytes from a given collection fail to fertilize (Yanagida, 2004) occur in 1–5% of all ICSI treatment cycles (Liu *et al.*, 1995). In such incidences of ICSI failure, the main cause has been shown to be the lack of oocyte activation (Javed *et al.*, 2010; Vanden Meerschaut *et al.*, 2013a; Neri *et al.*, 2014). The options for couples facing total fertilization failure are limited (Yuzpe *et al.*, 2000; Heindryckx *et al.*, 2005). Total fertilization failure cases are relatively rare, partly because typically ~10 oocytes are available for sperm injection (Sunkara *et al.*, 2011). This multiplicity of oocytes masks the fact that fertilization after ICSI is less effective when rated per oocyte injected. For

example, using sperm and oocytes with apparently normal parameters in cases where tubal blockage was the only evident cause of infertility, only 67% of oocytes were activated after sperm injection (Bukulmez *et al.*, 2000), implying that one in three sperm is ineffective in activating the oocyte. This recurrent failure per oocyte becomes a particular concern when oocytes are limited, in mild ovarian stimulation or natural ovulation cycles (Pelinck *et al.*, 2002; Loutradis *et al.*, 2007).

The activation of mammalian oocytes at fertilization involves an extensive series of Ca²⁺ transients, known as Ca²⁺ oscillations. Each Ca²⁺ spike lasts about 1 min and the Ca²⁺ transients occur at intervals of 5–30 min (Miyazaki and Ito, 2006; Swann and Lai, 2013). Such Ca²⁺ oscillations have been observed in human and mouse oocytes after *in vitro* fertilization and ICSI (Sato *et al.*, 1999). Such Ca²⁺ oscillations are both necessary and sufficient for oocyte activation (Kurokawa *et al.*, 2004) and may influence subsequent embryo development (Ducibella *et al.*, 2002; Yu *et al.*, 2008). Accumulating scientific and clinical evidence favours the idea that Ca²⁺ oscillations are triggered after sperm oocyte

membrane fusion, which allows the entry of a sperm-specific PLC isoform, termed PLC ζ , into the oocyte cytoplasm (Saunders et al., 2002; Nomikos et al., 2005). Injection of PLC ζ cRNA or recombinant protein stimulates parthenogenetic activation of mouse and human oocytes, and subsequent development of mouse embryos up to the blastocyst stage at rates similar to those seen after fertilization (Miyazaki, 1995).

Mouse oocytes have been frequently used as a model system for studying human IVF and ICSI. The mouse oocyte has also been used to directly test the ability of human sperm to cause oocyte activation or Ca²⁺ oscillations after cross-species ICSI (Araki et al., 2004; Vanden Meerschaut et al., 2013a). Injection of human sperm into mouse oocytes can trigger oocyte activation, but sperm from failed clinical ICSI cycles generally show a reduced ability to activate mouse oocytes after injection (Araki et al., 2004; Vanden Meerschaut et al., 2013a). Furthermore, injecting human sperm into mouse oocytes also causes a series of Ca²⁺ oscillations. The frequency of these human sperm-induced Ca²⁺ oscillations is often higher than observed at fertilization in mouse oocytes (Yoon et al., 2012; Nikiforaki et al., 2014). This is probably because human PLC ζ has a greater intrinsic potency than mouse PLC ζ and is able to cause the same pattern of Ca²⁺ oscillations at ~10 times lower concentrations than mouse PLC ζ (Yu et al., 2008). It has been reported that sperm from men with repeated fertilization failure, or with mutations in PLC ζ protein, show a markedly reduced ability to cause Ca²⁺ oscillations after ICSI into mouse oocytes (Yoon et al., 2008; Heytens et al., 2009; Kashir et al., 2011, 2012). However, it was noteworthy that even for men who have had successful ICSI cycles, only ~55% of normal sperm cause high-frequency Ca²⁺ oscillations in mouse oocytes (Vanden Meerschaut et al., 2013a; Nikiforaki et al., 2014). These data suggest that in cases of human ICSI failure, there might be either a deficiency, or relative lack of Ca²⁺ oscillations.

In cases of failed or poor rates of fertilization after ICSI, the only available treatment option is the use of artificial oocyte activation agents. This usually consists of treating oocytes with Ca²⁺ ionophores, such as A23187 or ionomycin, a procedure that has been successfully used to overcome fertilization failure in many cases (Kyono et al., 2012). Despite its apparent utility there are few controlled studies on the efficacy of Ca²⁺ ionophores as a means of rescuing failed oocyte activation and development. Ca²⁺ ionophore application, as currently used in most clinics, only causes a single large Ca²⁺ increase that does not mimic the series of Ca²⁺ oscillations seen at fertilization (Rinaudo et al., 1997). Stimuli that elicit multiple Ca²⁺ transients are known to be a more effective means of activating mammalian oocytes (Ozil and Swann, 1995; Ducibella et al., 2002). Sr²⁺ media can be used to cause such repetitive Ca²⁺ oscillations in mouse oocytes, but Sr²⁺ has not been shown to trigger Ca²⁺ release in human oocytes (Rogers et al., 2004). PLC ζ remains the only physiological agent that has been repeatedly shown to produce a prolonged series of Ca²⁺ oscillations in all mammalian oocytes studied, including human oocytes (Ito et al., 2011; Nomikos et al., 2013a, b; Kashir et al., 2014).

In this study, we have investigated whether recombinant human PLC ζ protein can be used to rescue cases of failed fertilization or to improve poor development rates after ICSI. We use a mouse model of failed oocyte activation after ICSI by applying mild heat treatment of mouse sperm. We find that heat-treated sperm display a reduced ability to generate Ca²⁺ oscillations in mouse oocytes, consistent with consequent observations of reduced preimplantation development. We also show

that fertilization failure and poor embryo development after ICSI with heat-treated sperm can be rescued by subsequent microinjection of recombinant human PLC ζ protein, which along with Sr²⁺ media, is more effective than Ca²⁺ ionophore. Moreover, we demonstrate that microinjection of PLC ζ after standard ICSI with normal, untreated sperm does not impair embryo development. These data provide the basis for future studies to examine the potential use of recombinant human PLC ζ as a biological therapeutic to rescue human oocytes from failed fertilization after ICSI.

Materials and Methods

All chemicals were obtained from Sigma Aldrich UK unless stated otherwise and were of embryo grade where available. M2 media was used for handling oocytes outside the incubator. KSOM was obtained from Merk-Millipore (Watford, UK), or made as previously described (Summers et al., 1995). The microinjection buffer was 100 mM KCl in 20 mM Hepes, pH 7.2 (Swann, 1990).

Gametes and embryos

Mouse oocytes were obtained from superovulated female MF1 mice as described previously (Yu et al., 2008). Cumulus-oocyte masses from oviducts were incubated in hyaluronidase in M2 media, and then isolated oocytes were washed in M2 media alone. Mouse embryos were cultured in KSOM media (Summers et al., 1995) under mineral oil in an incubator at 37°C gassed with 5% CO₂. Mouse sperm were collected from the cauda epididymis of euthanized male mice (C57xCBA F1 hybrid), released into T6 media (Jones et al., 1995) and in some cases, sperm were frozen in the same media (without cryopreservation) in a -80°C freezer. For heat treatment, sperm were incubated for the specified times in a water bath at 56°C then sonicated for ~10 s before being added to drops containing the oocytes. For Sr²⁺ activation after ICSI as described by Yoshida and Perry (2007) using heat inactivated sperm for 30 min, oocytes were allowed to recover for 30 min in M2. Oocytes were then incubated afterwards in calcium-free HKSOM supplemented with 5 mM Sr²⁺ for 2 h after which they are washed in M2 then cultured in KSOM under 5% CO₂ in air at 37°C.

Ethical approval

All procedures using animals were approved by Cardiff University Animals Ethics Committee and carried out under a UK Home Office Project Licence.

Microinjection

Mouse oocytes were microinjected with sperm using custom-made ICSI pipettes (Yoshida and Perry, 2007). The micropipettes containing the sperm were advanced through the oocyte plasma membrane using a piezo-pulse delivered by a Prime Tech piezo manipulation system (Intracel, Royston, UK). Recombinant PLC ζ was injected using a fine tip micropipette that was inserted into the oocyte using pressure pulses (Picopump, World Precision Instruments, USA), as described previously (Saunders et al., 2002).

Ca²⁺ measurements

Cytosolic free Ca²⁺ concentrations were measured in individual mouse oocytes by monitoring the fluorescence of PE3 or Rhod dextran (Takahashi et al., 1999). The PE3 dye was loaded into oocytes by incubating for 30 min in 10 μ M of PE3-AM (also known as fura2LeakRes-AM) dissolved in M2 media. PE3 is a similar dye to fura2 except that it does not undergo compartmentalization or extrusion from the cytosol and hence can be used for longer-term recordings of Ca²⁺ (Takahashi et al., 1999). Oocytes were then washed free of the AM dye and allowed to equilibrate for another ~30 min before making

fluorescence measurements. Rhod dextran was injected into oocytes before the start of experiments. The measurements of fluorescence were made by placing oocytes in drops of HKSOM media on a heated stage of a Nikon TiU epifluorescence microscope. For PE3 based recordings, oocytes were exposed to fluorescence excitation at 380 nm and emission measured at 510–550 nm (Takahashi *et al.*, 1999). The fluorescence ratio of the signals at 350 and 380 nm was then plotted against time as a measure of relative cytosolic Ca²⁺ concentrations. In a similar manner for Rhod dextran, the fluorescence was plotted against time with excitation at 550 nm and emission collected at 580–620 nm (Swann and Lai, 2013).

Recombinant protein expression and purification

Human PLC ζ was expressed as a NusA-6xHis-tagged fusion protein and purified by affinity chromatography on Ni-NTA resin using standard procedures (Qiagen) and elution with 275 mM imidazole, as described previously (Nomikos *et al.*, 2013b). NusA tagged human PLC ζ was tested for its ability to cause Ca²⁺ oscillations in oocytes prior to use and was injected at a pipette concentration of 0.01 mg/ml (Nomikos *et al.*, 2013a, b; Nikiforaki *et al.*, 2014). Control NusA protein was purified in a similar manner and when injected separately was used at a pipette concentration of 0.1 mg/ml in KCl/Hepes buffer.

Statistical analysis

Data on the number of Ca²⁺ transients and rate of embryo development from control and experimental groups were analysed using ANOVA or pairwise Student's *t*-tests. Differences at *P* < 0.05 were considered significant.

Results

Sperm treatment and Ca²⁺ oscillations after ICSI


To create a clinical scenario mimicking poor fertilization after ICSI, we exposed mouse sperm to heat treatment since previous studies have shown that the oocyte activating sperm factor is sensitive to mild heat treatment (Perry *et al.*, 1999). We incubated mouse sperm at 56°C for time intervals varying from 2.5 to 30 min. Figure 1A–C shows the recordings of Ca²⁺ oscillations observed in mouse oocytes after ICSI with either fresh (control) sperm or with sperm that had been heat treated for various times such as 20 or 30 min. ICSI using fresh sperm results in a series of Ca²⁺ oscillations during the first 2 h. In contrast, any heat treatment of sperm resulted in a significantly decreased number of Ca²⁺ spikes over the 2 h recording period. ICSI with sperm that had been heat treated for shorter durations such as 2.5 min produced some Ca²⁺ spikes (Fig. 1D), but significantly fewer than with control sperm. ICSI with frozen and thawed sperm caused slightly fewer Ca²⁺ oscillations in the 2 h recording period than freshly prepared sperm (Fig. 1D). Figure 1D also shows that the mean number of Ca²⁺ spikes is progressively reduced as the heat treatment is increased from 2.5 min, with a complete loss of Ca²⁺ oscillations after ICSI seen with a 30 min treatment. These data suggest that exposure of sperm at 56°C for increasing time periods is associated with a progressive reduction in the ability of sperm to trigger Ca²⁺ oscillations.

We further examined the effect of heat-treated ICSI sperm upon the oocyte activation rate and subsequent development to the blastocyst stage. We used MFI oocytes for this work where control rates of development *in vitro* to the blastocyst stage after fertilization are around 50%

(Ozil and Swann, 1995; Ducibella *et al.*, 2002). Figure 2 and Supplementary Table S1 show that the rate of oocyte activation, as defined by pronuclear formation, was progressively reduced by heat treating sperm for 2.5–30 min. Similarly, embryonic development to blastocysts was also reduced in embryos resulting from ICSI with heat-treated sperm. Figure 2 indicates that an increase in time of sperm exposure to 56°C is consistent with a progressive decrease in the number of ICSI embryos that develop to the 2-cell or blastocyst stage. These data suggest that mild heat treatment of mouse sperm leads to loss in the capacity of sperm to support both Ca²⁺ oscillations and preimplantation development after ICSI. The observed thermal impairment of sperm function mimics a clinical scenario of oocyte activation failure and poor development after IVF/ICSI.

The most common treatment for failed fertilization after ICSI is to expose oocytes to Ca²⁺ ionophores (Nasr-Esfahani *et al.*, 2010). Figure 3A shows that oocytes injected with sperm heat-treated for 30 min responded with a single large Ca²⁺ increase after exposure to ionomycin. For Fig. 3A the ionomycin was washed out after 5 min in order to mimic the duration of the Ca²⁺ transient seen in human oocytes exposed to ionophore (Rinaudo *et al.*, 1997). We then wanted to compare these stimuli to those that are known to cause Ca²⁺ oscillations. First we used Sr²⁺ media which, as shown in Fig. 3B, causes a series of Ca²⁺ oscillations (mean of 7.91 Ca²⁺ spikes in 2 h, with a standard deviation of 2.64, *n* = 15) after ICSI in a manner previously described (Perry *et al.*, 1999). However, in order to generate the Ca²⁺ oscillations in a way that would be effective in human oocytes, we used recombinant PLC ζ protein. After ICSI with 30 min heat-treated sperm, oocyte Ca²⁺ oscillations could be rescued by microinjection of recombinant human PLC ζ (Fig. 3C). The amount of PLC ζ protein we used is comparable to that used in previous experiments and was designed to generate a pattern of Ca²⁺ oscillations similar to those seen after ICSI with control sperm. Importantly, we also performed double injection experiments by injecting PLC ζ protein into oocytes that had been through ICSI with control (fresh) sperm (Fig. 4A). For these double-injected oocytes, there were also Ca²⁺ oscillations after PLC ζ injection, although we cannot distinguish the relative contribution to these oscillations from the PLC ζ injection and the injected sperm. It was, however, notable that the frequency of Ca²⁺ oscillations in these experiments was similar to that in control ICSI, or with ICSI using frozen–thawed sperm (see Fig. 4B).

The above experiments show that Ca²⁺ signals could be restored by the use of ionomycin, Sr²⁺ media, or by introducing recombinant PLC ζ protein, in oocytes injected with heat-treated sperm. We monitored oocyte activation and subsequent development in such oocytes. Figure 5 shows that following ICSI with sperm treated for 30 min and then incubated with Ca²⁺ ionophore, oocytes could be activated but only 13.3% formed blastocysts (see also Supplementary Table S2). In contrast, when we used the same procedure but replaced ionophore activation with Sr²⁺ media, or PLC ζ injection, not only were oocytes activated but more than 34–36% developed to the blastocysts stage. This is significantly more than with ionophore treatment. We found that the use of frozen sperm was less effective than fresh sperm in triggering development up to the blastocyst stage, but that injection of PLC ζ could effectively rescue these oocytes and give rise to developmental rates similar to those of fresh sperm (Supplementary Table S2). These data suggest that agents that induce Ca²⁺ oscillations, such as PLC ζ protein injection, provide an effective means of rescuing oocyte activation failure or



poor development after ICSI. We also noted that injecting either PLC ζ protein after ICSI using control sperm also led to oocyte activation and that the rate of 2-cell and blastocyst formation was similar to that seen with control sperm ICSI alone (Fig. 5 and Supplementary Table S2). These data show that the injection of PLC ζ , in addition to a normal Ca^{2+} oscillations stimulus from a control sperm, does not appear to impair preimplantation development to the blastocyst stage.

Discussion

Fertilization failure in an oocyte after ICSI occurs both sporadically in many treatment cycles and can sometimes occur in all oocytes. Whilst there may be many reasons for fertilization failure, the most common cause is failure of oocyte activation (Javed et al., 2010; Nasr-Esfahani

et al., 2010). In this study, we showed that mild heat treatment of mouse sperm leads to a reduction in its ability to trigger Ca^{2+} oscillations in mouse oocytes after ICSI. It is known that exposing mouse sperm to temperatures of $\sim 90^\circ\text{C}$ results in DNA damage, but that heat treatment at lower temperatures (56°C) does not cause DNA damage nor impair embryo development when an independent oocyte activation stimulus is used (Perry et al., 1999). Our present data confirm that sperm contain a heat labile factor that triggers Ca^{2+} oscillations and this sperm-borne factor is sensitive to relatively mild heat treatment of 56°C for 30 min. This heat treatment is similar to that previously used to inactivate SOAF (sperm oocyte activating factor) in earlier reports of mouse ICSI (Perry et al., 1999). Previous ICSI studies also demonstrated that the SOAF, which is present in the sperm perinuclear theca, is PLC ζ (Perry et al., 1999), as identified by Saunders et al. (2002). PLC ζ is presently




Figure 2 The percentage of embryo developmental to the blastocyst stage after ICSI is compared with that after ICSI using sperm that had been heat inactivated for various times. A significant difference for development to blastocyst was seen from control ICSI for the 10, 20, 30 min heat treatments (Student's *t*-test pairwise between the control and each heat treatment, $P = 0.0038$, 0.0023 , and 0.000 , respectively). Error bars are SEMs.

the only sperm-specific protein confirmed to have an intrinsic ability to cause Ca^{2+} oscillations in mammalian oocytes by multiple independent research teams (Saunders *et al.*, 2002; Kurokawa *et al.*, 2004; Rogers *et al.*, 2004; Miyazaki and Ito, 2006; Nomikos *et al.*, 2014). Therefore, it is probable that heat treatment leads to sperm PLC ζ inactivation in our experiments. Interestingly, freeze thawing of sperm reduces the activity and distribution of PLC ζ (Kashir *et al.*, 2011). This is consistent with our finding that this step slightly reduces the number of Ca^{2+} oscillations after ICSI as well as the percentage of embryos developing to the blastocyst stage. This could be due to some PLC ζ enzyme inactivation due to the freeze thaw cycle, or to loss of PLC ζ from the sperm. Nevertheless, in our study, defining the exact cause of loss of Ca^{2+} oscillations is not critical since we have identified a simple way of mimicking the clinical observation in humans where sperm from subfertile males shows a reduced ability to cause Ca^{2+} oscillations in oocytes after ICSI.

One advantage of using 56°C heat treatment to deplete the ability of sperm to activate oocytes after ICSI is that it enables provision of different time points that creates a range of effects. Indeed we found that varying the duration of sperm heating from 2.5 to 30 min led to progressive loss of Ca^{2+} oscillation-inducing activity. Along with the loss of Ca^{2+} oscillations, there was a loss of oocyte activation and embryo development up to the blastocyst stage. We found that embryo development to the blastocyst stage fell from $\sim 50\%$ down to 0% correlating directly with the reduced number of Ca^{2+} spikes caused by increasingly longer heat treatments. It should be noted that we used mouse oocytes from the MFI strain which develop to blastocysts *in vitro* at a rate of $\sim 50\%$,






Figure 3 Intracellular Ca^{2+} measured in mouse oocytes treated with artificial stimuli after ICSI. Conditions are the same as in Fig. 1 except that PE3 was used to record Ca^{2+} in (B) and (C) where fluorescence is in arbitrary (ratio) units (a.u.). In (A) is shown an example of oocyte treated with ionomycin for 5 min ($n = 21$ recordings) where the bar indicates the time for which ionomycin was applied. In (B) is an example of an oocyte injected with heat treatment (30 min) inactivated sperm and then treated with Sr^{2+} media ($n = 12$). In (C) is shown an example of an oocyte injected with heat treatment (30 min) inactivated sperm and then subsequently injected with purified recombinant PLC ζ protein ($n = 15$).

which matches clinical scenarios with human embryos. These data suggests that an attenuation of the Ca^{2+} oscillations following ICSI leads to decreased rates of oocyte activation and development. This is a



scenario that appears to occur in some cases of subfertility found by practitioners of ICSI (Vanden Meerschaut et al., 2013a; Nikiforaki et al., 2014).

In clinical cases where poor ICSI oocyte activation rates are observed, the current treatment option is the use of Ca^{2+} ionophores, which provide an artificial stimulus (Neri et al., 2014). It has been used in many cases of complete activation failure. It can give rise to live births in treatment cycles that would otherwise fail (Yanagida et al., 2008; Nasr-Esfahani et al., 2010), but the efficiency of this activation protocol compared with a physiological stimulus has not been critically examined. Ca^{2+} ionophores can only cause one or, in some protocols, two large Ca^{2+} increases, which fails to mimic the multiple Ca^{2+} oscillations that occur at fertilization (Vincent et al., 1992; Heytens et al., 2008). In our study, we tested the ability of ionomycin to overcome failed oocyte activation after ICSI with heat-treated sperm and found that it could



restore activation. Ionomycin treatment also lead to development to the blastocyst stage, but it was less effective than either Sr^{2+} media or recombinant PLC ζ at producing oocyte activations. PLC ζ recombinant protein has been previously shown to overcome activation failure after ICSI in human oocytes (Yoon et al., 2012). This set a precedent for the use of PLC ζ as a means of rescuing oocytes from activation failure.

However, the efficiency of activation was not compared with other protocols. In our current experiments, the higher rates of Sr²⁺ and PLC ζ -mediated oocyte activation versus ionophore-mediated oocyte activation also lead to better rates of embryo development to the blastocyst. Even though it may not be usable for human oocytes, we used Sr²⁺ media because it induces Ca²⁺ oscillations. Consequently our data indicated that it is because PLC ζ causes Ca²⁺ oscillations that it is a more effective stimulus for rescuing oocyte activation after ICSI failure than Ca²⁺ ionophores. This is consistent with previous data showing that multiple Ca²⁺ transients are the most effective way to activate mouse or rabbit oocytes (Ducibella *et al.*, 2002, 2006).

In our protocol, we have used a second injection to introduce PLC ζ and induced Ca²⁺ oscillations in the mouse oocyte after ICSI. The use of a second injection has been used in at least one clinic where Ca²⁺ is injected after ICSI, and then ionophores are applied, as part of a protocol to rescue oocyte from ICSI failure (Vanden Meerschaut *et al.*, 2013a, b). Human oocytes are more robust than mouse oocytes since mouse oocytes specifically require piezo devices, or electrical oscillation, for pipette insertion (Swann, 1990; Yoshida and Perry, 2007). Hence, a second injection of PLC ζ protein should be possible in a clinical setting. We additionally anticipate that introducing PLC ζ into an oocyte should be relatively safe for embryos. So far the treatment of oocytes with Ca²⁺ ionophores, which produces an unnaturally large rise in Ca²⁺, has been shown to have no discernible damaging effect on embryos or live young (Yanagida, 2004; Vanden Meerschaut *et al.*, 2013a). PLC ζ causes Ca²⁺ oscillations indistinguishable from those seen at fertilization and so should not lead to any effects upon development other than those arising naturally.

Previous studies have suggested that human sperm contain variable amounts of PLC ζ and also have a variable ability to cause Ca²⁺ oscillations in oocytes (Grasa *et al.*, 2008; Kashir *et al.*, 2013). During ICSI, an embryologist is unable to assess the ability of a chosen sperm to cause Ca²⁺ oscillations. If PLC ζ is injected into ICSI oocytes to overcome cases of suspected activation failure, then it is possible that these oocytes will receive some endogenous Ca²⁺ releasing activity from the sperm in addition to that from the injected PLC ζ . We tested the developmental consequence of this by examining the scenario where control sperm was used for ICSI and then the same oocytes were subsequently injected with the single sperm equivalent dose of PLC ζ used for rescuing complete activation failure. We also injected PLC ζ into oocytes that had ICSI with frozen–thawed sperm and have only a slightly reduced number of Ca²⁺ oscillations. In both these cases of ICSI + PLC ζ injection, we found that after oocyte activation, the development rates to the blastocyst stage were similar to that for control ICSI. This benign effect might seem surprising since previous studies have shown that PLC ζ over-expression can lead to poor development to the blastocyst stage (Yu *et al.*, 2008). However, the previous deleterious effects on development required over-expression of about 10-fold more PLC ζ than is effective in causing oocyte activation. In the current study, we estimate that oocytes should have received little more than double the normal PLC ζ level in one sperm. The lack of change in the frequency of Ca²⁺ oscillations is consistent with previous studies in mouse oocytes that have found only a marginal difference in Ca²⁺ oscillation frequency after fertilization by two versus one sperm in zona-free oocytes (Faure *et al.*, 1999). These data suggest that PLC ζ can be injected at a dose to successfully activate an oocyte that had not been activated by ICSI, and that this will not lead to detrimental effects on early embryo development.

Supplementary data

Supplementary data are available at <http://molehr.oxfordjournals.org/>

Acknowledgements

We thank Junaid Kashir for helpful comments on the manuscript.

Authors' roles

R.S. and Y.Y. carried out experiments on oocytes and embryos and analysed the data. M.N. purified and characterized the recombinant human PLC ζ protein. K.S. and F.A.L. planned and supervised the project. All authors prepared the manuscript.

Funding

R.S. is supported by a research scholarship from the Libyan Ministry of Education. M.N. holds an EU-FP7 Marie Curie Intra-European Fellowship (628634). This work was also partly funded by a research grant from Cook Medical Technologies LLC. Funding to pay the Open Access publication charges for this article was provided by Cardiff University.

Conflict of interest

F.A.L. and K.S. hold patents with Cardiff University on PLC ζ Sperm factor sequences. Other authors declare no conflict of interest.

References

- Araki Y, Yoshizawa M, Abe H, Murase Y. Use of mouse oocytes to evaluate the ability of human sperm to activate oocytes after failure of activation by intracytoplasmic sperm injection. *Zygote* 2004;**12**:111–116.
- Bukulmez O, Yarali H, Yucel A, Sari T, Gurgan T. Intracytoplasmic sperm injection versus in vitro fertilization for patients with a tubal factor as their sole cause of infertility: a prospective, randomized trial. *Fertil Steril* 2000;**73**:38–42.
- Ducibella T, Huneau D, Angelichio E, Xu Z, Schultz RM, Kopf GS, Fissore R, Madoux S, Ozil JP. Egg-to-embryo transition is driven by differential responses to Ca²⁺ oscillation number. *Dev Biol* 2002;**250**:280–291.
- Ducibella T, Schultz RM, Ozil JP. Role of calcium signals in early development. *Semin Cell Dev Biol* 2006;**17**:324–332.
- Faure JE, Myles DG, Primakoff P. The frequency of calcium oscillations in mouse eggs at fertilization is modulated by the number of fused sperm. *Dev Biol* 1999;**213**:370–377.
- Grasa P, Coward K, Young C, Parrington J. The pattern of localization of the putative oocyte activation factor, phospholipase C ζ , in uncapacitated, capacitated, and ionophore-treated human spermatozoa. *Hum Reprod* 2008;**23**:2513–2522.
- Heindryckx B, Van Der Elst J, De sutter P, Dhont M. Treatment option for sperm- or oocyte-related fertilization failure: assisted oocyte activation following diagnostic heterologous ICSI. *Hum Reprod* 2005;**20**:2237–2241.
- Heytens E, Soleimani R, Lierman S, De Meester S, Gerris J, Dhont M, Van der Elst J, De Sutter P. Effect of ionomycin on oocyte activation and embryo development in mouse. *Reprod Biomed Online* 2008;**17**:764–771.
- Heytens E, Parrington J, Coward K, Young C, Lambrecht S, Yoon SY, Fissore RA, Hamer R, Deane CM, Ruas M *et al.* Reduced amounts and abnormal forms of phospholipase C zeta (PLC ζ) in spermatozoa from infertile men. *Hum Reprod* 2009;**24**:2417–2428.

- Ito J, Parrington J, Fissore RA. PLC ζ and its role as a trigger of development in vertebrates. *Mol Reprod Dev* 2011; **78**:846–853.
- Javed M, Esfandiari N, Casper RF. Failed fertilization after clinical intracytoplasmic sperm injection. *Reprod Biomed Online* 2010; **20**:56–67.
- Johnson LNC, Sasson IE, Sammel MD, Dokras A. Does intracytoplasmic sperm injection improve the fertilization rate and decrease the total fertilization failure rate in couples with well-defined unexplained infertility? A systematic review and meta-analysis. *Fertil Steril* 2013; **100**:704–711.
- Jones KT, Carroll J, Merriman JA, Whittingham DG, Kono T. Repetitive sperm-induced Ca²⁺ transients in mouse oocytes are cell cycle dependent. *Development* 1995; **121**:3259–3266.
- Kashir J, Jones C, Lee HC, Rietdorf K, Nikiforaki D, Durran C, Ruas M, Tee ST, Heindryckx B, Galione A et al. Loss of activity mutations in phospholipase C zeta (PLC ζ) abolishes calcium oscillatory ability of human recombinant protein in mouse oocytes. *Hum Reprod* 2011; **26**:3372–3387.
- Kashir J, Konstantinidis M, Jones C, Lemmon B, Chang Lee H, Hamer R, Heindryckx B, Deane CM, De Sutter P, Fissore RA et al. A maternally inherited autosomal point mutation in human phospholipase C zeta (PLC ζ) leads to male infertility. *Hum Reprod* 2012; **27**:222–231.
- Kashir J, Jones C, Mounce G, Ramadan WM, Lemmon B, Heindryckx B, De Sutter P, Parrington J, Turner K, Child T et al. Variance in total levels of phospholipase C zeta (PLC- ζ) in human sperm may limit the applicability of quantitative immunofluorescent analysis as a diagnostic indicator of oocyte activation capability. *Fertil Steril* 2013; **99**:107–117.e3.
- Kashir J, Nomikos M, Lai FA, Swann K. Sperm-induced Ca²⁺ release during egg activation in mammals. *Biochem Biophys Res Commun* 2014; **450**:1204–1211.
- Kurokawa M, Sato KI, Fissore RA. Mammalian fertilization: from sperm factor to phospholipase C ζ . *Biol Cell* 2004; **96**:37–45.
- Kyono K, Takisawa T, Nakajo Y, Doshida M, Toya M. Birth and follow-up of babies born following ICSI with oocyte activation using strontium chloride or calcium ionophore A23187. *J Mamm Ova Res* 2012; **29**:35–40.
- Liu J, Nagy Z, Joris H, Tournaye H, Smits J, Camus M, Devroey P, Van Steirteghem A. Analysis of 76 total fertilization failure cycles out of 2732 intracytoplasmic sperm injection cycles. *Hum Reprod* 1995; **10**:2630–2636.
- Loutradis D, Drakakis P, Vomvolaki E, Antsaklis A. Different ovarian stimulation protocols for women with diminished ovarian reserve. *J Assist Reprod Genet* 2007; **24**:597–611.
- Miyazaki S. Calcium signalling during mammalian fertilization. *Ciba Found Symp* 1995; **188**:235–247; discussion 247.
- Miyazaki S, Ito M. Calcium signals for egg activation in mammals. *J Pharmacol Sci* 2006; **100**:545–552.
- Nasr-esfahani MH, Deemeh MR, Tavalae M. Artificial oocyte activation and intracytoplasmic sperm injection. *Fertil Steril* 2010; **94**:520–526.
- Neri QV, Lee B, Rosenwaks Z, Machaca K, Palermo GD. Understanding fertilization through intracytoplasmic sperm injection (ICSI). *Cell Calcium* 2014; **55**:24–37.
- Nikiforaki D, Vanden Meerschaut F, De Gheselle S, Qian C, Van den Abbeel E, De vos WH, Deroo T, De Sutter P, Heindryckx B. Sperm involved in recurrent partial hydatidiform moles cannot induce the normal pattern of calcium oscillations. *Fertil Steril* 2014; **102**:581–588.e1.
- Nomikos M, Blayney LM, Larman MG, Campbell K, Rossbach A, Saunders CM, Swann K, Lai FA. Role of phospholipase C- ζ domains in Ca²⁺-dependent phosphatidylinositol 4,5-bisphosphate hydrolysis and cytoplasmic Ca²⁺ oscillations. *J Biol Chem* 2005; **280**:31011–31018.
- Nomikos M, Kashir J, Swann K, Lai FA. Sperm PLC ζ : From structure to Ca²⁺ oscillations, egg activation and therapeutic potential. *FEBS Lett* 2013a; **587**:3609–3616.
- Nomikos M, Yu Y, Elgmati K, Theodorido M, Campbell K, Vassilakopoulou V, Zikos C, Livanou E, Amso N, Nounesis G et al. Phospholipase C ζ rescues failed oocyte activation in a prototype of male factor infertility. *Fertil Steril* 2013b; **1**:76–85.
- Nomikos M, Sanders JR, Theodoridou M, Kashir J, Matthews E, Nounesis G, Lai FA, Swann K. Sperm-specific post-acrosomal WW-domain binding protein (PAWP) does not cause Ca²⁺ release in mouse oocytes. *Mol Hum Reprod* 2014; **20**:938–947.
- Ozil JP, Swann K. Stimulation of repetitive calcium transients in mouse eggs. *J Physiol* 1995; **483**:331–346.
- Palermo G, Joris H, Devroey P, Van Steirteghem AC. Pregnancies after intracytoplasmic injection of single spermatozoon into an oocyte. *Lancet* 1992; **340**:17–18.
- Pelinc M, Hoek A, Simons AHM, Heineman MJ. Efficacy of natural cycle IVF: a review of the literature. *Hum Reprod Update* 2002; **8**:129–139.
- Perry ACF, Wakayama T, Yanagimachi R. A novel trans-complementation assay suggests full mammalian oocyte activation is coordinately initiated by multiple, submembrane sperm components. *Biol Reprod* 1999; **60**:747–755.
- Rinaudo P, Massobrio M, Pepperell JR, Keefe DL, Buradgunta S. Dissociation between intracellular calcium elevation and development of human oocytes treated with calcium ionophore. *Fertil Steril* 1997; **68**:1086–1092.
- Rogers NT, Hobson E, Pickering S, Lai FA, Braude P, Swann K. Phospholipase C ζ causes Ca²⁺ oscillations and parthenogenetic activation of human oocytes. *Reproduction* 2004; **128**:697–702.
- Sato MS, Yoshitomo M, Mohri T, Miyazaki S. Spatiotemporal analysis of [Ca²⁺]_i rises in mouse eggs after intracytoplasmic sperm injection (ICSI). *Cell Calcium* 1999; **26**:49–58.
- Saunders CM, Larman MG, Parrington J, Cox LJ, Royse J, Blayney LM, Swann K, Lai FA. PLC ζ : a sperm-specific trigger of Ca²⁺ oscillations in eggs and embryo development. *Development* 2002; **129**:3533–3544.
- Summers MC, Bhatnagar PR, Lawitts JA, Biggers JD. Fertilization in vitro of mouse ova from inbred and outbred strains: complete preimplantation embryo development in glucose-supplemented KSOM. *Biol Reprod* 1995; **53**:431–437.
- Sunkara SK, Rittenberg V, Raine-fenning N, Bhattacharya S, Zamora J, Coomarasamy A. Association between the number of eggs and live birth in IVF treatment: an analysis of 400 135 treatment cycles. *Hum Reprod* 2011; **26**:1768–1774.
- Swann K. A cytosolic sperm factor stimulates repetitive calcium increases and mimics fertilization in hamster eggs. *Development* 1990; **110**:1295–1302.
- Swann K, Lai FA. PLC ζ and the initiation of Ca²⁺ oscillations in fertilizing mammalian eggs. *Cell Calcium* 2013; **53**:55–62.
- Takahashi A, Camacho P, Lechleiter JD, Herman B. Measurement of intracellular calcium. *Physiol Rev* 1999; **79**:1089–1125.
- Vanden Meerschaut F, Leybaert L, Nikiforaki D, Qian C, Heindryckx B, De Sutter P. Diagnostic and prognostic value of calcium oscillatory pattern analysis for patients with ICSI fertilization failure. *Hum Reprod* 2013a; **28**:87–98.
- Vanden Meerschaut F, Nikiforaki D, De Roo C, Lierman S, Qian C, Schmitt-john T, De sutter P, Heindryckx B. Comparison of pre-and post-implantation development following the application of three artificial activating stimuli in a mouse model with round-headed sperm cells deficient for oocyte activation. *Hum Reprod* 2013b; **28**:1190–1198.
- Vincent C, Cheek TR, Johnson MH. Cell cycle progression of parthenogenetically activated mouse oocytes to interphase is dependent on the level of internal calcium. *J Cell Sci* 1992; **103**:389–396.
- Yanagida K. Complete fertilization failure in ICSI. *Hum Cell* 2004; **17**:187–193.
- Yanagida K, Fujikura Y, Katayose H. The present status of artificial oocyte activation in assisted reproductive technology. *Reprod Med Biol* 2008; **7**:133–142.
- Yoon SY, Jellerette T, Salicioni AM, Hoi CL, Yoo MS, Coward K, Parrington J, Grow D, Cibelli JB, Visconti PE et al. Human sperm devoid of PLC, zeta 1 fail to induce Ca²⁺ release and are unable to initiate the first step of embryo development. *J Clin Invest* 2008; **118**:3671–3681.

- Yoon SY, Eum JH, Lee JE, Lee HC, Kim YS, Han JE, Won HJ, Park SH, Shim SH, Lee WS et al. Recombinant human phospholipase C zeta 1 induces intracellular calcium oscillations and oocyte activation in mouse and human oocytes. *Hum Reprod* 2012;**27**:1768–1780.
- Yoshida N, Perry ACF. Piezo-actuated mouse intracytoplasmic sperm injection (ICSI). *Nat Protoc* 2007;**2**:296–304.
- Yu Y, Saunders CM, Lai FA, Swann K. Preimplantation development of mouse oocytes activated by different levels of human phospholipase C zeta. *Hum Reprod* 2008;**23**:365–373.
- Yuzpe AA, Liu Z, Fluker MR. Rescue intracytoplasmic sperm injection (ICSI)—salvaging in vitro fertilization (IVF) cycles after total or near-total fertilization failure. *Fertil Steril* 2000;**73**:1115–1119.