

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <http://orca.cf.ac.uk/96491/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

El-Hiti, Gamal A., Smith, Keith, Hegazy, Amany S., Masmali, Ali M. and Kariuki, Benson 2016. Crystal structure of 3-tert-butyl-7-azadioxindole, C₁₁H₁₄N₂O₂. Zeitschrift für Kristallographie - New Crystal Structures 231 (4) , pp. 1069-1070. 10.1515/ncrs-2016-0054 file

Publishers page: <http://dx.doi.org/10.1515/ncrs-2016-0054> <<http://dx.doi.org/10.1515/ncrs-2016-0054>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Gamal A. El-Hiti*, Keith Smith, Amany S. Hegazy, Saud A. Alanazi and Benson M. Kariuki

Crystal structure of 2-(4-methoxyphenyl)-1,3-thiazolo[4,5-*b*]pyridine, C₁₃H₁₀N₂OS




Table 1: Data collection and handling.

Crystal:	Colourless plate Size 0.42 × 0.11 × 0.03 mm
Wavelength:	Mo K α radiation (0.71073 Å)
μ :	2.7 cm ⁻¹
Diffractometer, scan mode:	SuperNova, ω
2 θ _{max} , completeness:	59.4°, >99%
$N(hkl)$ _{measured} , $N(hkl)$ _{unique}	6077, 6077
Criterion for I_{obs} , $N(hkl)$ _{gt} :	$I_{obs} > 2 \sigma(I_{obs})$, 3490
$N(param)$ _{refined} :	309
Programs:	SHELX [5], CrysAlis [6], WinGX [7]

DOI 10.1515/ncrs-2016-0053

Received February 12, 2016; accepted June 4, 2016; available online June 15, 2016

Abstract

C₁₃H₁₀N₂OS, monoclinic, $P2_1/c$ (no 14), $a = 14.182(3)$ Å, $b = 5.9674(6)$ Å, $c = 26.683(4)$ Å, $\beta = 101.162(17)^\circ$, $V = 2215.4(6)$ Å³, $Z = 8$, $R_{gt}(F) = 0.0634$, $wR_{ref}(F^2) = 0.1514$, $T = 150$ K.

CCDC no.: 1483462

The asymmetric unit of the title crystal structure is shown in the figure. Tables 1 and 2 contain details of the measurement method and a list of the atoms including atomic coordinates and displacement parameters.

Source of material

2-(4-Methoxyphenyl)-1,3-thiazolo[4,5-*b*]pyridine was obtained in 64% yield from reaction of 4-anisic acid and 3-(diisopropylaminothiocarbonylthio)-2-aminopyridine in

*Corresponding author: Gamal A. El-Hiti, Cornea Research Chair, Department of Optometry, College of Applied Medical Sciences, King Saud University, P.O. Box 10219, Riyadh 11433, Saudi Arabia, e-mail: gelhiti@ksu.edu.sa

Keith Smith, Amany S. Hegazy and Benson M. Kariuki: School of Chemistry, Cardiff University, Main Building, Park Place, Cardiff CF10 3AT, UK

Saud A. Alanazi: Cornea Research Chair, Department of Optometry, College of Applied Medical Sciences, King Saud University, P.O. Box 10219, Riyadh 11433, Saudi Arabia

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	<i>x</i>	<i>y</i>	<i>z</i>	U_{iso}^*/U_{eq}
C1	0.6236(3)	0.4406(7)	0.01361(17)	0.0228(10)
C2	0.6729(3)	0.6468(7)	0.09315(16)	0.0253(10)
C3	0.6364(3)	0.4282(7)	0.09712(16)	0.0204(10)
C4	0.7035(3)	0.7725(7)	0.13689(16)	0.0310(10)
H4	0.7271	0.9210	0.1356	0.037*
C5	0.6977(3)	0.6689(8)	0.18229(17)	0.0342(11)
H5	0.7192	0.7468	0.2135	0.041*
C6	0.6614(3)	0.4541(8)	0.18367(16)	0.0307(11)
H6	0.6585	0.3907	0.2160	0.037*
C7	0.6060(3)	0.3708(6)	-0.04020(16)	0.0178(9)
C8	0.5598(3)	0.1690(7)	-0.05484(15)	0.0228(10)
H8	0.5396	0.0786	-0.0296	0.027*
C9	0.5423(3)	0.0955(7)	-0.10473(15)	0.0219(10)
H9	0.5087	-0.0408	-0.1139	0.026*
C10	0.5746(3)	0.2244(7)	-0.14126(15)	0.0244(10)
C11	0.6216(3)	0.4273(7)	-0.12777(16)	0.0244(10)
H11	0.6431	0.5154	-0.1530	0.029*
C12	0.6368(3)	0.4994(7)	-0.07809(15)	0.0245(10)
H12	0.6685	0.6381	-0.0692	0.029*
C13	0.5125(4)	-0.0308(7)	-0.20808(16)	0.0359(12)
H13A	0.5449	-0.1593	-0.1893	0.054*
H13B	0.5095	-0.0523	-0.2448	0.054*
H13C	0.4472	-0.0179	-0.2013	0.054*
C14	0.8828(3)	-0.0573(7)	0.00793(16)	0.0210(9)
C15	0.8865(3)	0.1647(7)	0.08535(15)	0.0240(10)
C16	0.9264(3)	-0.0497(8)	0.09164(17)	0.0258(10)
C17	0.8857(3)	0.2996(8)	0.12715(15)	0.0284(10)
H17	0.8584	0.4454	0.1235	0.034*
C18	0.9263(3)	0.2147(8)	0.17452(16)	0.0317(10)
H18	0.9288	0.3016	0.2046	0.038*

Table 2 (continued)

Atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso} */ <i>U</i> _{eq}
C19	0.9631(3)	−0.0009(8)	0.17702(18)	0.0375(12)
H19	0.9890	−0.0578	0.2101	0.045*
C20	0.8632(3)	−0.1351(7)	−0.04439(16)	0.0216(10)
C21	0.9034(3)	−0.3353(7)	−0.05740(15)	0.0228(10)
H21	0.9439	−0.4184	−0.0314	0.027*
C22	0.8863(3)	−0.4163(7)	−0.10669(16)	0.0252(10)
H22	0.9145	−0.5530	−0.1146	0.030*
C23	0.8270(3)	−0.2945(8)	−0.14471(16)	0.0249(10)
C24	0.7868(3)	−0.0925(7)	−0.13265(16)	0.0258(10)
H24	0.7465	−0.0087	−0.1586	0.031*
C25	0.8057(3)	−0.0157(7)	−0.08326(15)	0.0246(10)
H25	0.7788	0.1228	−0.0755	0.030*
C26	0.8395(4)	−0.5675(8)	−0.20821(17)	0.0379(12)
H26A	0.9099	−0.5647	−0.2010	0.057*
H26B	0.8162	−0.5964	−0.2447	0.057*
H26C	0.8169	−0.6863	−0.1881	0.057*
N1	0.6086(2)	0.3174(6)	0.05099(13)	0.0236(8)
N2	0.6299(3)	0.3306(6)	0.14154(13)	0.0289(9)
N3	0.9245(2)	−0.1732(6)	0.04762(13)	0.0239(8)
N4	0.9657(3)	−0.1362(6)	0.13766(14)	0.0335(10)
O1	0.5651(2)	0.1696(5)	−0.19173(10)	0.0309(7)
O2	0.8038(2)	−0.3560(5)	−0.19494(10)	0.0298(7)
S1	0.67055(8)	0.7093(2)	0.03003(4)	0.0279(3)
S2	0.84480(8)	0.2144(2)	0.02155(4)	0.0270(3)

phosphorus oxychloride under reflux for 4 h [1] or in 82% yield from treatment of 3-(diisopropylaminothiocarbonylthio)-2-(4-methoxybenzoylamino)pyridine with hydrochloric acid under reflux for 5 h [2]. Colourless crystals of the title compound were obtained *via* crystallization from diethyl ether.

Experimental details

The crystal was twinned and refinement was performed using the option HKLF 5 in SHELX [5]. H atoms were placed in calculated positions and refined using a riding model, with *U*_{iso}(H) set to 1.2*U*_{eq}(C) and aromatic C–H distance of 0.93 Å. The methyl (C–H = 0.96 Å) groups were allowed to rotate around the C–C bond (HFIX 137 option in SHELX [5]) with *U*_{iso}(H) set to 1.5*U*_{eq}(C).

Discussion

Various thiazolopyridines have been synthesised *via* various synthetic methods and many derivatives have shown interesting biological applications [3, 4]. The asymmetric unit of the crystal structure of the title compound consists of two independent molecules of C₁₃H₁₀N₂OS. The methoxyphenyl groups in both molecules are planar, as illustrated by C_{Me}-O-C_{Ph}-C_{Ph} torsion angles of 3.8(5)° and 3.4(6)°. The dihedral angles between the planes through the methoxyphenyl and thiazolopyridine groups of the two molecules are 9.6(2)° and 10.0(1)°, indicating that both molecules are nearly planar. In the crystal, π-π interaction occurs between inversion-related molecules separated by a distance of 3.4 Å of the corresponding planes.

Acknowledgements: The authors extend their appreciation to the College of Applied Medical Sciences Research Center and the Deanship of Scientific Research at King Saud University for their funding of this research.

References

- El-Hiti, G. A.: A convenient procedure for the formation of 2-substituted thiazolopyridines. *Monatsh. Chem.* **134** (2003) 837–841.
- Smith, K.; Anderson, D.; Matthews, I.: A convenient synthesis of 2-substituted thiazolo[4,5-*b*]pyridines via directed metalation. *Sulfur Lett.* **18** (1995) 79–95.
- Cai, D.; Z., Z.-H.; Chen, Y.; Yan, X.-J.; Zou, L.-J.; Wang, Y. X.; Liu, X.-Q.: Synthesis, antibacterial and antitubercular activities of some 5*H*-thiazolo[3,2-*a*]pyrimidin-5-ones and sulfonic acid derivatives. *Molecules* **20** (2015) 16419–16434.
- Thomae, D.; Perspicace, E.; Hesse, S.; Kirsch, G.; Seck, P.: Synthesis of substituted [1, 3]thiazolo[4,5-*b*]pyridines and [1, 3]thiazolo[4,5-*d*][1, 2, 3]triazines. *Tetrahedron* **64** (2008) 9309–9314.
- Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr.* **A64** (2008) 112–122.
- Agilent. CrysAlis PRO. Agilent technologies, Yarnton, England, 2014.
- Farrugia, L. J.: WinGX and ORTEP for Windows: an update. *J. Appl. Crystallogr.* **45** (2012) 849–854.