

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository:<https://orca.cardiff.ac.uk/id/eprint/109938/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Sanislav, I. V., Dirks, P. H. G. M., Blenkinsop, Thomas and Kolling, S. L. 2018. The tectonic history of a crustal-scale shear zone in the Tanzania Craton from the Geita Greenstone Belt, NW-Tanzania Craton. *Precambrian Research* 310 , pp. 1-16.  
10.1016/j.precamres.2018.02.025

Publishers page: <http://dx.doi.org/10.1016/j.precamres.2018.02.025>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



# 1      **The tectonic history of a crustal-scale shear zone in the Tanzania**

## 2                      **Craton from the Geita Greenstone Belt, NW-Tanzania Craton**

3      **I. V. Sanislav<sup>1\*</sup>, P. H. G. M. Dirks<sup>1</sup>, T. Blenkinsop<sup>2</sup>, S. L. Kolling<sup>3</sup>**

4      *<sup>1</sup>Economic Geology Research Centre (EGRU) and Geoscience Department, James Cook*  
5      *University, Townsville, 4011, QLD, Australia; e-mail: [ioan.sanislav@jcu.edu.au](mailto:ioan.sanislav@jcu.edu.au); phone:*  
6      *(+61) 07 4781 3293; fax: (+61) 07 4781 5581*

7      *<sup>2</sup>School of Earth & Ocean Sciences, Cardiff University, Cardiff CF10 3AT, United Kingdom*

8      *<sup>3</sup>Geita Gold Mine, Geita, P.O. Box 532, Geita Region, Tanzania*

### 9 10      **Abstract**

11              In this contribution, we present for the first-time field based evidence of a crustal scale  
12      shear zone from the southern margin of Geita Greenstone Belt. The Geita Shear Zone is a broad  
13      (~ 800 m wide) ductile, high-strain, deformation zone that can be traced for at least 50 km  
14      along the southern margin of the Geita Greenstone Belt. It is near vertical, trends ~ E-W and  
15      separates the mafic volcanics of the Kiziba Formation, to the north, from the TTG gneisses that  
16      crop out south of the shear zone. The shear zone is hosted almost entirely by the TTG gneisses  
17      and is characterised by a well-developed mylonitic foliation near the greenstone margin that  
18      transitions into a gneissic foliation and eventually becomes a weakly developed foliation  
19      further south. It contains approximately equal amounts of dextral, sinistral, and asymmetric  
20      shear sense indicators suggesting that the shear zone accommodated mainly flattening strain  
21      while the mineral stretching lineation defined by quartz and feldspar ribbons and stretched  
22      biotite selvages plunges shallowly W. A series of younger, sub-vertical, NW trending brittle-

23 ductile, strike-slip shear zones truncate and displace the Geita Shear Zone with dextral  
24 displacement in the order of 2-4 km. Deformed tonalite interpreted to predate the shear zone  
25 yielded U-Pb zircon ages of ~2710 Ma while synshearing granodiorite samples have zircon  
26 ages between 2680 Ma and 2660 Ma. The ~2630 Ma age of the undeformed Nyankumbu  
27 granite is interpreted to mark the minimum age of movement on the shear zone. The presence  
28 of 3000 Ma and 3200 Ma zircon xenocrysts in the tonalite and granodiorite opens the possibility  
29 that older basement rocks underlie the greenstone belts in the northern half of Tanzania Craton.  
30 Whether or not the greenstone belts were erupted on older basement, thrust on top of older  
31 basement rocks or incorporated older basement fragments has profound implications for the  
32 tectonic framework and evolution of the Tanzania Craton.

33

## 34 **1. Introduction**

35

36 Large scale shear zones are well documented features in Archean terrains worldwide  
37 (e.g. Bédard et al., 2003; Cassidy et al., 2006; Chardon et al., 2008; Jelsma and Dirks, 2002;  
38 Dirks et al., 2013) and their structural style has implications for the tectonic history and crustal  
39 growth models of Archean cratons. One of the most common Archean structural styles is the  
40 dome and keel geometry (e.g. Pilbara, Kaapvaal, Western Dharwar), linked to gravity-driven  
41 tectonics and redistribution of rock masses through vertical processes (e.g. Choukroune et al.,  
42 1995; Chardon et al., 1996; Collins et al., 1998; Bédard et al., 2003; Van Kranendonk, 2011).  
43 The majority of these types of structures are characterised by gneiss domes with radial  
44 elongation lineations and normal shear sense indicators suggesting diapiric ascent through the  
45 crust. Between the gneiss domes, synformal-shaped greenstone belts are preserved. In general,  
46 the geometry of the greenstone belts follows the contours of the gneiss domes resulting in cusp-

47 like shapes. Typical examples include the Barberton Greenstone Belt in South Africa (e.g.  
48 Dirks et al., 2013; Van Kranendonk et al., 2014; Brown, 2015) and the greenstone belts in the  
49 Pilbara Craton of Western Australia (e.g. Collins et al., 1998; Van Kranendonk et al., 2004).  
50 Alternative interpretations for dome and keel geometries include core complexes exposed by  
51 extensional unroofing (Kloppenburg et al., 2001) or antiformal culminations exposed by cross-  
52 folding (Blewett et al., 2004).

53 Another common Archean structural style is represented by linear belts separated by  
54 transcurrent shear zones, which are typically interpreted to indicate lateral terrain accretion  
55 (e.g. Polat et al., 1998; Dirks et al., 2002; Blewett et al., 2010; Kabete et al., 2012a). Typical  
56 examples include the Yilgarn Craton (e.g. Cassidy et al., 2006; Blewett et al., 2010) the  
57 Superior Province (e.g. Card, 1990; Polat et al., 1998) and the Dharwar Craton (e.g. Chadwick  
58 et al., 2000; Manikyamba and Kerrich, 2012). The transcurrent shear zones form an  
59 anastomosing pattern encompassing elongated greenstone belts, many with similar  
60 stratigraphy, bordered by granite-gneiss terranes of younger age. For example, in the eastern  
61 Yilgarn Craton, detailed stratigraphic analyses showed that the stratigraphy of the greenstone  
62 belts, although disrupted by shear zones, can be correlated for hundreds of kilometres (Hayman  
63 et al., 2015).

64 The structural and tectonic evolution of the Archean Tanzania Craton is poorly  
65 understood (Kabete et al., 2012a; Kabete et al., 2012b) and in many instances stratigraphic  
66 correlations based on sparse geochronological data are contradictory (e.g. Borg and Krogh,  
67 1999; Many et al., 2006; Sanislav et al., 2014). A first attempt to define tectonic and structural  
68 boundaries within the Tanzania Craton was made by Kabete et al. (2012a) based on existing  
69 geological maps and the interpretation of geophysical datasets. A series of NW trending shear  
70 zones were proposed and interpreted to delineate superterrane boundaries (Fig. 1). However,  
71 the existence of these large-scale shear zones, their continuity, style, age and kinematic sense

72 was never confirmed with field studies, and the terrane-accretion model, therefore, remains  
73 entirely speculative.

74 In this contribution, we provide the first field-based description of a regional scale shear  
75 zone from the northern half of the Tanzania Craton and present new zircon age data from  
76 igneous rocks that intruded along and across the shear zone to constrain its kinematic history.  
77 Results will be discussed in terms of the regional significance for the structural and tectonic  
78 evolution of this part of the Tanzania Craton.

79

## 80 **2. Regional geology**

81

82 The geology of the Archean Tanzania Craton is generally described in terms of three  
83 main stratigraphic and tectonic units, the Dodoman Supergroup, The Nyanzian Supergroup and  
84 the Kavirondian Supergroup (e.g. Many and Maboko, 2008; Kabete et al., 2012a; Cook et al.,  
85 2016; Sanislav et al. 2017). The Dodoman Supergroup is the oldest tectonic unit and its  
86 occurrence is restricted mainly to the central part of the Tanzania Craton (Kabete et al., 2012a).  
87 This unit consists of high-grade mafic and felsic granulite with subordinate lower-grade schist  
88 and thin slivers of greenstone. A limited number of zircon ages confirm that the Dodoman  
89 Supergroup forms the older part of the Tanzania Craton with ages ranging between 3 Ga and  
90 3.6 Ga (Kabete et al., 2012b).

91 The Nyanzian and the Kavirondian Supergroups are concentrated in the northern half  
92 of the Tanzania Craton and were placed stratigraphically above the Dodoman Supergroup (e.g.  
93 Quenell et al 1956; Gabert, 1990). The Nyanzian Supergroup is overlain unconformably  
94 (Gabert, 1990) by the Kavirondian Supergroup. The lowermost unit of the Nyanzian is

95 dominated by mafic volcanics and volcanoclastics with minor felsic volcanics erupted at ~  
96 2820 Ma (e.g. Borg and Krogh, 1999; Many and Maboko, 2003; Cook et al., 2016). These are  
97 overlain by felsic volcanic and pyroclastic units inter-bedded with banded ironstone,  
98 volcanoclastic sequences and immature turbiditic sediments (e.g. Borg, 1992; Borg and  
99 Shackleton, 1997; Sanislav et al., 2015). All greenstone sequences from the northern half of  
100 Tanzania were assigned to the Nyanzian Supergroup. Six major greenstone belts were  
101 described (Fig. 1): the Sukumaland Greenstone Belt (SGB), The Nzega Greenstone Belt, the  
102 Iramba-Sekenke Greenstone Belt, the Shynianga-Malita Greenstone Belt, the Kilimafedha  
103 Greenstone Belt and the Musoma-Mara Greenstone Belt (Borg and Shackleton, 1997). The  
104 greenstone belts are fragmented by shear zones and intruded by ~2800 to 2600 Ma granitoids  
105 (Kabete et al., 2012b; Sanislav et al., 2014). Most of the greenstone belts are oriented along ~  
106 E-W lineaments (Fig. 1), one of these lineaments forming the southern margin of the Geita  
107 Greenstone Belt. The Kavirondian Supergroup consists mainly of coarse-grained  
108 conglomerate, grit and quartzite (Gabert, 1990) and its areal extent is poorly constrained.

109         The Geita Greenstone Belt (GGB) is a large greenstone fragment forming most of the  
110 northern part of what has previously been defined as the SGB (Fig. 1). It contains elements of  
111 the Nyanzian and the Kavirondian stratigraphy (Fig. 2). The base of the greenstone belt is  
112 dominated by lower-amphibolite facies metamorphosed mafic volcanics forming the Kiziba  
113 Formation (Cook et al., 2016). Whole rock Sm-Nd dating of the mafic volcanics yield model  
114 ages of ~2820 Ma (Many and Maboko, 2008; Cook et al., 2016). The mafic volcanics of the  
115 Kiziba Formation are positioned below a complexly deformed sequence of intercalated  
116 volcanoclastics, ironstones and immature turbidites, which experienced greenschist facies  
117 metamorphism (Cook et al., 2016; Sanislav et al., 2015, 2017). The greenstone sequence was  
118 intruded syntectonically by diorite (D<sub>2</sub> to D<sub>4</sub>) and by late-syntectonic felsic porphyries (D<sub>6</sub> to  
119 D<sub>7</sub>). The northern, eastern and western margins of the greenstone are intruded by 2660 to 2620

120 Ma high-K granites (Sanislav et al., 2014). The southern margin of the Geita Greenstone Belt  
121 is bounded by gneiss and deformed granite across a steeply dipping shear zone, which was  
122 intruded by the undeformed high-K Nyankumbu granite (Fig. 2). A few outcrops of  
123 Kavirondian dominated by quartzitic conglomerate with minor sandstone occur in the north-  
124 eastern part of the greenstone belt.

125 Sanislav et al. (2015; 2017) proposed that the GGB was affected by at least eight,  
126 Archean deformation events. The first event ( $D_1$ ) produced mainly layer parallel shears in what  
127 may have been an extensional setting, while the second ( $D_2$ ) to fourth ( $D_4$ ) deformation events  
128 were responsible for the formation of large-scale folds and associated shear zones, which in  
129 turn were refolded by open to gentle, sub-horizontal ( $D_5$ ) folds. These ductile events were  
130 followed by two brittle-ductile shearing events that produced localised reverse faults ( $D_6$ ) and  
131 large scale NW trending dextral strike slip faults ( $D_7$ ). The last event ( $D_8$ ) is an extensional  
132 event that produced steeply dipping and commonly ~ E-W trending normal faults that  
133 reactivated earlier  $D_6$ - $D_7$  structures.

134

### 135 **3. Geita Shear Zone**

136

137 The Geita Shear Zone (GSZ) is a mylonitic shear zone that defines the southern margin  
138 of the GGB and can be mapped along strike for at least 50 km (Fig. 2). The general orientation  
139 of the GSZ is ~ E-W (Fig. 2) and the overall straight map pattern and gravity data for the  
140 greenstone belt suggest that the shear zone is steeply dipping. Regional aeromagnetic data  
141 suggest that the GSZ extends further to the east and west (Fig. 1) outside of the area mapped  
142 in Figure 2. In the Geita area the shear zone marks the boundary between mafic volcanics of  
143 the Kiziba Formation to the north and TTG granitoid-dominated gneiss and granites to the

144 south (Fig. 2). The shear zone is generally poorly exposed, and is characterised on  
145 aeromagnetic data sets by a 700-900 m wide zone with an intensely developed planar fabric  
146 (Fig. 2 ), and in the field by scattered outcrops of partly recrystallised mylonite and striped  
147 gneiss (Fig. 3a and b) Aeromagnetic datasets further indicate the presence of well-foliated  
148 gneiss across a width of at least 0.8 km to the south of the main shear zone, whilst to the north,  
149 the shear zone appears to transect well-foliated meta basalt with the northern boundary of the  
150 GSZ defined by a low-angle truncation zone within deformed meta-basalt. The contact between  
151 the Kiziba Formation and the TTG gneisses is not exposed anywhere in the Geita Greenstone  
152 Belt, but the two units crop out within 50 meters of each other SW of Geita Town (GR 411520-  
153 9679010 - all grid references in WGS84-36S), which represents one of the few areas with  
154 excellent outcrop in the shear zone.

155         Near the contact with the Kiziba Formation the TTG gneiss units preserve a mylonitic  
156 fabric characterised by an intense, mm-scale foliation and a generally near-horizontal mineral  
157 stretching lineation defined by quartz-feldspar rodding (Fig. 3a). The mylonitic fabric has been  
158 recrystallised, resulting in recovery of the deformation textures on thin-section scale, and  
159 partial destruction of the mineral lineation especially in compositionally homogenous (i.e. bt-  
160 amp-poor) portions of the gneiss. Highly boudinaged quartz veins and flattened feldspar  
161 porphyroclasts (Fig. 3c) with symmetrical and asymmetrical tails (Figs. 3d and e) are locally  
162 preserved within the mylonitic fabric, and shear bands are common some containing granite  
163 veins. Going south, away from the contact with the greenstone belt, the mylonitic fabric  
164 remains strong for about 400m beyond which the deformation fabric becomes less intense and  
165 takes the form of a well-developed gneissic foliation with alternating dark biotite selvages and  
166 cm-scale quartzo-feldspathic lithons that retain the stretching lineation (Fig. 3f). At about 1 km  
167 south of the contact only a weak gneissic foliation is preserved (Fig. 3g) and feldspar

168 phenocrysts appear largely undeformed (Fig. 3h and i). The shear fabric is truncated by the  
169 undeformed Nyankumbu granite (Fig. 3j).

170 The TTG gneiss units and granite intrusions south of the GSZ display a single dominant  
171 gneissic foliation that is steep with a dominantly southerly dip and an E-W trend (average dip  
172 / dip direction:  $86^{\circ}/170^{\circ}$ ; Fig. 4a). The stretching lineation in the gneisses and granites  
173 generally plunges shallowly W (Fig. 4b). Plagioclase porphyroclasts are mostly symmetrical,  
174 but some porphyroclasts with asymmetric tails indicate variable shear directions with dextral  
175 (Fig. 3d) appearing to be more common on a regional scale. Dextral and sinistral shear bands  
176 occur side-by side. This suggests that the shear zone mainly accommodated flattening strain  
177 with N-S shortening, and stretch along a horizontal, E-W direction, possibly with a minor  
178 dextral component.

179 In the metabasalts to the north of the contact the GSZ is not exposed but geophysical  
180 images indicate that near the contact with the TTG the fabric from the Kiziba Formation is  
181 truncated and deflected into parallelism with the GSZ suggesting that the shear zone has only  
182 a limited extent to the north. Rare metabasalt outcrops preserve evidence of intrafolial folding  
183 (Fig. 6a; Cook et al., 2016) and a foliation defined by the preferential alignment of amphibole  
184 and locally by a well-defined mm-scale banding of alternating layers of amphibole and  
185 plagioclase. The foliation trends ~E-W and dips mainly north (average dip / dip direction  
186  $73^{\circ}/352^{\circ}$ ; Fig. 4c). These mafic schists locally contain a moderately E to NE plunging (Fig.  
187 4d) mineral lineation defined by hornblende and a plagioclase stretching lineation (Fig. 2), and  
188 numerous asymmetric, boudinaged quartz veins (Fig. 6b) indicative of a N-up, sinistral  
189 movement sense. Aeromagnetic images suggest that locally, large-scale folds occur within the  
190 Kiziba formation north of the GSZ with E-W trending fold axes and axial planar surfaces (Fig.  
191 3g). In the western part of the GGB, within metabasalts that outcrop to the north of the GSZ,

192 flattened pillow basalts occur with well-preserved vesicles filled with quartz and carbonate  
193 indicating younging to the north (Cook et al., 2016).

194         Along the entire length of the GGB, the GSZ has been truncated and displaced by NW-  
195 trending, near-vertical, D<sub>7</sub> faults (Fig. 2; Sanislav et al., 2015; 2017). These faults contain near-  
196 horizontal lineations defined by quartz fibres and striations and brittle-ductile deformation  
197 textures including well preserved S-C fabrics indicative of a dextral sense of shear. The D<sub>7</sub>  
198 shear zones are characterised by numerous, planar milky quartz veins that intruded parallel to  
199 the strike of the D<sub>7</sub> faults. Networks of conjugate pairs of NW- and NE-trending quartz veins  
200 are also common in the mylonitic gneisses of the GSZ within the vicinity of D<sub>7</sub> faults. Dextral  
201 displacements of the GSZ along the principle D<sub>7</sub> faults are in the order of 2-4 km.

202         Within the felsic gneiss along the Geita Shear zone, tonalitic to granodioritic gneiss is  
203 the dominant rock type, but less deformed, porphyritic to coarse-grained granite and  
204 granodiorite bodies (Figs. 3h and i) are common as dykes and small intrusions. In the zone of  
205 most intense deformation a number of over-printing dyke-like (Fig. 3b) to lensoidal bodies of  
206 granite and granodiorite can be recognised, most of which are positioned in near-parallelism to  
207 the main gneissic fabric and all of which contain an S-L fabric with constant lineation  
208 directions. This suggests that shearing coincided with the emplacement of several generations  
209 of felsic intrusions, which can provide a direct age estimate for the principle period of  
210 movement along the shear zone.

211         Away from the core of the shear zone, some of the truncating granite dykes show only  
212 weak fabrics or no deformation features at all. Some granite dykes did not develop a foliation,  
213 but preserve a stretching lineation parallel to the dominant lineation direction in the shear zone,  
214 defined by quartz aggregates, aligned biotite grains and weakly deformed feldspar phenocrysts.  
215 The TTG gneisses are medium- to coarse-grained and their mineralogy is dominated by quartz

216 and plagioclase with minor amounts of k-feldspar, biotite and hornblende. The cross cutting  
217 granite dykes and bodies intruding the gneisses of the GSZ during the shearing events are  
218 quartz-rich and have approximately equal amounts of plagioclase and k-feldspar with only  
219 minor amounts of biotite and hornblende. Based on their mineralogy the granites have been  
220 classified as monzogranites. It is common for these granites to contain large k-feldspar  
221 phenocrysts (Figs. 3h and i).

222 A younger, undeformed granite, the Nyankumbu granite (Fig. 3j) intruded the GSZ  
223 (Fig. 2) in the central southern part of the greenstone belt, and crosscuts the mylonitic fabrics  
224 in the TTG gneisses and the Kiziba Formation. This granite provides a lower limit for the time  
225 of shearing along the GSZ. The mineralogy of the Nyankumbu granite is dominated by quartz  
226 and k-feldspar with lesser plagioclase, biotite and minor hornblende, and it contains numerous  
227 xenoliths. The xenoliths vary in size from a few centimetres to tens of centimetres in diameter  
228 and their composition is similar to the composition of the TTG gneiss units and granites that  
229 are intruded by the Nyankumbu granite.

230

#### 231 **4. Geochronology**

232

233 Nine samples were collected for zircon dating along and across the GSZ (Fig. 2) to  
234 constrain the time of shearing. Seven samples come from the granitoids south of the GSZ and  
235 two samples (K81 and IH3) come from small tonalite bodies that intruded into the Kiziba  
236 Formation. These tonalite bodies are part of the greenstone sequence affected by the GSZ and  
237 their time of emplacement is interpreted to provide an upper age limit for the GSZ. From south  
238 of the GSZ two samples were collected from the TTG gneisses (SM02 and SM25), which may  
239 provide a maximum time for shearing. Three samples were collected from the deformed

240 granites that intruded into the TTG gneisses and that retain the shear fabric, i.e. these granites  
241 may have been intruded during shearing (SM07, SM23 and SM27). Two samples were  
242 collected from the undeformed Nyankumbu granite (SM56 and SM57) that crosscuts the GSZ  
243 and constrain the cessation of shearing along the GSZ.

244 All samples were dated at the Advanced Analytical Centre hosted by James Cook  
245 University. All zircon grains were imaged with a Jeol JSM5410LV with attached  
246 cathodoluminescence detector. Analytical spots for each zircon grain were selected based on  
247 internal zonation. U-Pb isotope analyses were obtained using a GeoLas 200 Excimer Laser  
248 Ablation System in a He ablation atmosphere, coupled to a Varian ICP-MS 820 series  
249 instrument. The detailed analytical procedures used for U-Pb dating in the James Cook  
250 University laboratory are outlined in Sanislav et al. (2014). Data were processed using the  
251 software package GLITTER<sup>TM</sup> (Jackson et al., 2004), and age calculations were done using  
252 Isoplot (Ludwig, 2003). The zircon standard GJ1 (ID-TIMS <sup>207</sup>Pb/<sup>206</sup>Pb age = 608.5 ± 0.4 Ma;  
253 Jackson et al., 2004) was used as the primary standard while the zircon standards FC1 (ID-  
254 TIMS <sup>207</sup>Pb/<sup>206</sup>Pb age = 1099.0 ± 0.6 Ma; Paces and Miller, 1993) and 91500 (<sup>207</sup>Pb/<sup>206</sup>Pb age  
255 = 1065 ± 0.3 Ma; Wiedenbeck et al., 1995) were used as secondary standards.

256 The weighted average ages (errors at 95% confidence) for the standards were as  
257 follows: GJ-1 - 610 ± 10 Ma for <sup>207</sup>Pb/<sup>206</sup>Pb and 600 ± 2 Ma for <sup>206</sup>Pb/<sup>238</sup>U; FC-1 - 1101 ± 9  
258 Ma for <sup>207</sup>Pb/<sup>206</sup>Pb and 1103 ± 15 Ma for <sup>206</sup>Pb/<sup>238</sup>U; 91500 - 1071 ± 17 Ma for <sup>207</sup>Pb/<sup>206</sup>Pb and  
259 1060 ± 5 Ma for <sup>206</sup>Pb/<sup>238</sup>U. Analyses with significant discordance (>10%) and those with  
260 elevated common Pb, where <sup>206</sup>Pb/<sup>204</sup>Pb (background corrected) is <1000, were excluded from  
261 age calculations.

262

## 263 5. Results

264

### 265 5.1. The TTG samples predating the GSZ

266

267 Samples SM02 and SM25 were collected from the TTG gneisses that crop out in the lower-  
268 strain, southern margin of the GSZ, where the shear fabric is only weakly developed. Both  
269 samples are coarse-grained and based on their mineralogy can be classified as tonalite. The  
270 zircon grains separated from these samples are, in general, euhedral to subhedral with a  
271 prismatic to needle-like shape and low cathodoluminescence luminosity (Fig. 6). The zircon  
272 grains separated from sample SM02 have a low luminescence core, without any clear zoning,  
273 surrounded by a thin rim overgrowth with a slightly higher luminescence (Fig. 6a). In some  
274 zircons the rim overgrowth is delineated from the core by a thin high luminescence band.  
275 Because the rim overgrowth was too thin to analyse by laser, all the measurements were  
276 performed on the cores. Eight U-Pb analyses with low  $^{204}\text{Pb}$  yielded concordant ages and were  
277 used for age calculations. The  $^{207}\text{Pb}/^{206}\text{Pb}$  weighted average of this sample is  $2705\pm 14$  Ma (Fig.  
278 6b) with an identical concordia age of  $2705\pm 18$  Ma (Fig. 6c), interpreted to represent the  
279 emplacement age.

280 The cathodoluminescence images of the zircon grains from sample SM25 revealed the  
281 presence of grains with complex internal growth domains (Fig. 6d). The majority of the zircons  
282 grains contain a low luminescence core with a weakly developed concentric zoning surrounded  
283 by a very thin, high luminescence rim (Fig. 6d). Two concordant and nine nearly concordant  
284 analyses from this domain have a  $^{207}\text{Pb}/^{206}\text{Pb}$  weighted average of  $2706\pm 13$  Ma (Fig. 6e) and  
285 an identical concordia age of  $2706\pm 13$  Ma (Fig. 6f), which are interpreted to represent the  
286 emplacement age. One large zircon grain (Fig. 6d) contains an elongated and sub-rounded

287 inherited core with low luminescence and without any internal zoning surrounded by a thick  
288 median overgrowth with higher luminescence and concentric zoning. The median overgrowth  
289 has a thin and weakly developed high luminosity rim. Two U-Pb analyses from the inherited  
290 core yielded a concordant analysis with a  $^{207}\text{Pb}/^{206}\text{Pb}$  age of  $2840\pm 22$  Ma and one nearly  
291 concordant analysis with a  $^{207}\text{Pb}/^{206}\text{Pb}$  age of  $2826\pm 21$  Ma. One nearly concordant analysis  
292 from the median overgrowth gave a  $^{207}\text{Pb}/^{206}\text{Pb}$  age of  $2710\pm 22$  Ma, which is similar to the  
293 emplacement age. Three zircon grains contained large, low luminescence cores surrounded by  
294 well-developed, high luminescence rims (Fig. 6d). Only one core analysis returned a nearly  
295 concordant  $^{207}\text{Pb}/^{206}\text{Pb}$  age of  $2702\pm 20$  Ma, similar to the emplacement age, while the rims  
296 returned nearly concordant analyses with a  $^{207}\text{Pb}/^{206}\text{Pb}$  weighted average age of  $2661\pm 24$  Ma  
297 (Fig. 6e) and a concordia age of  $2665\pm 41$  Ma (Fig. 6f). The rim age is interpreted to represent  
298 zircon overgrowth during a later thermal event related to the emplacement of the granites that  
299 intruded the TTG's during shearing along the GSZ.

300

### 301 *5.2. Tonalite intrusions internal to the greenstone belt predating the GSZ*

302

303 Samples IH3 and K81 were collected from two small tonalite bodies that intruded into the  
304 mafic rocks of the Kiziba Formation (Fig. 2). Both tonalite outcrops contain a well-developed  
305 foliation having a similar orientation to the foliation found in the surrounding mafic volcanics,  
306 which occur to the north of the GSZ. Sample IH3 was collected near the contact with the TTG  
307 gneisses where the foliation is sup-parallel to the trend of GSZ. Sample K81 was collected  
308 further north of the GSZ boundary where the foliation forms a small angle to the trend of the  
309 GSZ. The tonalite bodies are interpreted to have been emplaced before development of the  
310 GSZ. The zircon grains separated from these samples are generally medium- to small-sized,

311 prismatic, and mostly euhedral in shape. Cathodoluminescence images of zircon grains from  
312 sample IH3 (Fig. 7a) show that most of the zircon grains have low luminescence and poorly  
313 developed, concentric zoning. It is common for these zircon grains to display a thin, high-  
314 luminescence band, which only partly surrounds the grain, and which, in turn, is overgrown by  
315 a narrow rim (Fig. 7a). Ten concordant and near concordant analyses from these zircons gave  
316 a  $^{207}\text{Pb}/^{206}\text{Pb}$  weighted average age of  $2708\pm 15$  Ma (Fig. 7b) and a similar concordia age of  
317  $2708\pm 25$  Ma (Fig. 7c). These ages are interpreted to indicate the emplacement age of the  
318 tonalite bodies. The rim overgrowth was too narrow to be analysed. Sample IH3 also contained  
319 a few higher luminescence zircon grains (Fig. 7a) with some of them having a narrow rim  
320 overgrowth. Four analyses of these higher luminescence zircon grains returned concordant or  
321 near concordant ages. Their  $^{207}\text{Pb}/^{206}\text{Pb}$  weighted average age is  $2831\pm 21$  Ma (Fig. 7b) while  
322 the concordia age is  $2831\pm 22$  Ma (Fig. 7c). These ages are similar to the ages of the inherited  
323 zircon cores found in sample SM25 suggesting that these two intrusions may have sampled a  
324 similar reservoir.

325         Sample K81 yielded a large number of zircon grains. Cathodoluminescence images  
326 revealed that the majority of zircon grains are similar to the zircon grains separated from sample  
327 IH3; in general, they display low luminescence, with poorly developed concentric zoning, with  
328 or without a thin outer high-luminescence rim and a further, narrow rim overgrowth (Fig. 7d).  
329 Twenty-three concordant and nearly concordant analyses from these zircons gave a  $^{207}\text{Pb}/^{206}\text{Pb}$   
330 weighted average age of  $2711\pm 8$  Ma (Fig. 7e) and an identical concordia age of  $2711\pm 8$  Ma  
331 (Fig. 7f), which are interpreted to indicate the emplacement age. However, many of these  
332 zircon grains contain irregularly shaped, inherited cores with very low luminescence and no  
333 internal zoning (Fig. 7d). These cores always contain a thin and highly luminescent outer band  
334 that may or may not completely surround the core and have a narrow rim overgrowth. A few  
335 of these cores were large enough to be analysed. One nearly concordant analysis from a low

336 luminescence core yielded a  $^{207}\text{Pb}/^{206}\text{Pb}$  age of  $3231\pm 19$  Ma. Six other analyses performed on  
337 inherited cores yielded almost identical ages so that they were treated together. The  $^{207}\text{Pb}/^{206}\text{Pb}$   
338 weighted average age of these six analyses is  $3004\pm 21$  Ma (Fig. 7e) while their concordia age  
339 is  $3005\pm 26$  Ma (Fig. 7f).

340

### 341 *5.3. The granite samples emplaced during movement on the GSZ*

342

343 Three samples (SM07, SM23 and SM27) were collected from variably deformed,  
344 coarse-grained to porphyritic granite that intruded into the TTG gneisses south of the GSZ (Fig.  
345 2). They appear as dykes and lensoidal intrusions having a similar, but less intense shear fabric  
346 with the host TTG. Some thin dykes (~10 cm) of similar composition have been rotated into  
347 near parallelism with the shear fabric suggesting that these granites were emplaced during  
348 movement on the GSZ. The zircon grains are small- to medium-sized, in general prismatic and  
349 euhedral in shape (Figs. 8 and 9). They display low luminescence with a weakly defined growth  
350 zoning. All three samples contain zircon grains with inherited cores that are surrounded by a  
351 narrow high luminescence rim. However, only sample SM07 contained a zircon grain with an  
352 inherited core large enough to be analysed. This zircon grain (Fig. 8a) contains three growth  
353 domains: a large core surrounded by a very thin luminescent rim followed by a narrow median  
354 overgrowth, also surrounded by a thin high luminescent rim, and finally a rim overgrowth. One  
355 nearly concordant analysis from the core returned a  $^{207}\text{Pb}/^{206}\text{Pb}$  age of  $3264\pm 20$  Ma, which is  
356 similar to the age of one of the inherited cores found in sample K81. This sample also contained  
357 a zircon grain (Fig. 8a) with different cathodoluminescence properties compared to the  
358 remaining grains, including a well-defined growth zoning and medium luminescence. Two  
359 analyses were performed on this grain: one concordant analysis returned a  $^{207}\text{Pb}/^{206}\text{Pb}$  age of

360 2841±22 Ma and one nearly concordant analysis returned a  $^{207}\text{Pb}/^{206}\text{Pb}$  age of 2866±23 Ma.  
361 These ages are similar to the ages of inherited cores found in samples SM25 and IH3. Analyses  
362 performed on the remaining low luminescence grains yielded similar ages with a  $^{207}\text{Pb}/^{206}\text{Pb}$   
363 weighted average age of 2667±19 Ma (Fig. 8b) and a concordia age of 2667±27 Ma (Fig. 8c),  
364 which are interpreted to indicate emplacement ages. The zircon grains from samples SM23 and  
365 SM27 had small inherited cores, which could not be analysed. All zircon grains from sample  
366 SM23 displayed low luminescence (Fig. 8d) and yielded a  $^{207}\text{Pb}/^{206}\text{Pb}$  weighted average age of  
367 2671±18 Ma (Fig. 8e) and a concordia age of 2673±22 Ma (Fig. 8f). The zircon grains from  
368 sample SM27 have well defined growth zoning (Fig. 9a), and ten analyses yielded a  $^{207}\text{Pb}/^{206}\text{Pb}$   
369 weighted average age of 2680±13 Ma (Fig. 9b) and a concordia age of 2678±27 Ma (Fig. 9c).

370

#### 371 *5.4. The Nyankumbu granite emplaced into the GSZ*

372

373 Two samples (SM56 and SM57) were collected from the Nyankumbu granite to constrain  
374 the minimum age of the GSZ. Sample SM57 yielded a large number of zircon grains while  
375 sample SM56 yielded only few zircon grains. Both samples contain a mixed zircon population  
376 with inherited cores and xenocrysts. Cathodoluminescence images show that in sample SM56  
377 the majority of the grains are in general high luminescence with well-defined growth zoning  
378 (Fig. 10a). The inherited cores have low luminescence and in many instances are surrounded  
379 by rim overgrowths with a higher luminescence. A few zircon grains with low luminescence  
380 and a weakly defined growth zoning were interpreted as xenocrysts. Five high luminescence  
381 zircon grains yielded nearly concordant results with a  $^{207}\text{Pb}/^{206}\text{Pb}$  weighted average age of  
382 2638±16 Ma (Fig. 10b) and an identical concordia age (Fig. 10c). Two low luminescence  
383 zircon xenocrysts yielded a combined  $^{207}\text{Pb}/^{206}\text{Pb}$  age of 2655±25 Ma (Fig. 10b). Three zircon

384 grains contained inherited cores that were large enough to be analysed (Fig. 10a). Their  
385  $^{207}\text{Pb}/^{206}\text{Pb}$  ages are similar so that they were treated together and their  $^{207}\text{Pb}/^{206}\text{Pb}$  weighted  
386 average age is  $2709\pm 22$  Ma (Fig. 10b). The  $2638\pm 16$  Ma age is interpreted to indicate the  
387 emplacement age.

388 The zircon grains from sample SM57 have similar luminescence (Fig. 10d) and shape,  
389 which made it difficult to identify the zircon xenocrysts. Only a few grains contained inherited  
390 cores, but they were too small to be analysed. A number of 22 zircon grains from this sample  
391 returned concordant (15 grains) or nearly concordant (7 grains) ages. The youngest concordant  
392 grain returned a  $^{207}\text{Pb}/^{206}\text{Pb}$  age of  $2615\pm 21$  Ma while the oldest concordant grain returned a  
393  $^{207}\text{Pb}/^{206}\text{Pb}$  age of  $2684\pm 20$  Ma. The  $2615\pm 21$  Ma grain is similar within error to the weighted  
394 average of high luminescence grains from sample SM56, while the  $2684\pm 20$  Ma grain is similar  
395 within error to the ages of the deformed granites (Fig. 10e). Taking into account that the  
396 Nyankumbu granite intruded into the TTG and the deformed granites, we can assume that all  
397 grains with ages that are within error similar to the youngest age of the deformed granites or of  
398 the TTG's are xenocrysts, while all grains with ages that are within error similar to sample  
399 SM56 and younger than the syn-shearing granites belong to the Nyankumbu granite (Fig. 10e).  
400 Based on this assumption, 7 grains can be interpreted as Nyankumbu granite zircons and the  
401 remaining 15 grains can be interpreted as xenocrysts inherited from the surrounding older  
402 granites. The  $^{207}\text{Pb}/^{206}\text{Pb}$  weighted average age of the 7 zircon grains interpreted to be related  
403 to the emplacement of the Nyankumbu granite is  $2632\pm 16$  Ma (Fig. 10e) with a similar  
404 concordia age of  $2637\pm 22$  Ma (Fig. 10c). These ages are identical within error to the age of the  
405 high luminescence zircon grains from sample SM56, and indicate that the emplacement of the  
406 Nyankumbu granite took place between  $\sim 2632$  and  $2638$  Ma.

407

## 408        **6. Discussion**

409

### 410        *6.1. Structural history of the Geita Shear Zone*

411

412            The history of the GSZ can be reconstructed based on the field relationships and the  
413 new age data. Although, the contact between the TTG gneisses and the Kiziba Formation is not  
414 exposed, the map pattern (Fig. 2), suggests that the deformation fabric in the Kiziba Formation  
415 predates the deformation fabric in the TTG gneisses. For example, in the western part of the  
416 GGB (Fig. 2) the foliation in the Kiziba Formation is truncated by the GSZ while further east  
417 the foliation is deflected from NW trending into sub-parallelism with the foliation in the TTG,  
418 suggesting that at least near the shear zone margin the GSZ affected the Kiziba Formation.

419            The TTG gneisses and the deformed granites, south of the shear zone, and the mafic  
420 volcanics of the Kiziba Formation north of the shear zone, have similarly trending foliations  
421 (Figs. 4a and c), but differently oriented mineral lineations (Figs. 4b and d) and associated  
422 kinematic history, further supporting a different origin for the two deformation fabrics. The  
423 Kiziba Formation is characterised by moderately NE plunging lineations with top to the south  
424 shear-sense indicators, consistent with SW directed sinistral reverse movement while the  
425 shallowly W plunging lineations and the lack of a dominant shear sense direction along the  
426 mylonitic margin of the GSZ are consistent with N-S directed flattening strain. The kinematics  
427 of the Kiziba formation is similar to the main period of deformation of the greenstone belt,  
428 which produced the dominant D<sub>2</sub>-D<sub>3</sub> folding with SW-directed vergence and the sinistral-  
429 reverse D<sub>6</sub> shears present in the Geita Hill and Nyankanga deposits (Sanislav et al., 2015;  
430 2017). Both the GSZ and the Kiziba Formation are affected and displaced by a series of dextral-  
431 NW trending brittle-ductile shear zones (Fig. 2). These NW trending shear zones are of D<sub>7</sub>

432 origin (Sanislav et al., 2015; 2017) meaning that the GSZ must predate D<sub>7</sub> deformation. In the  
433 Geita Hill deposit, 2699 Ma (Borg and Krogh, 1999) diorite dykes are folded by the D<sub>3</sub> folds  
434 suggesting that the main period of folding occurred after 2700 Ma. The high-K granites that  
435 intruded the eastern, northern and western part of the greenstone belt between 2660 Ma and  
436 2620 Ma (Sanislav et al., 2014) truncate all the deformation fabric in the greenstone including  
437 the D<sub>7</sub> structures indicating a minimum age of 2660 Ma for the D<sub>7</sub> deformation. The ~ 2700  
438 Ma (Fig. 6) tonalite bodies that intruded the Kiziba Formation preserve the same deformation  
439 fabric with the host metabasalts indicating that the deformation of the Kiziba Formation  
440 occurred most probably after 2700 Ma. Taking into account the sinistral-reverse kinematics  
441 with top to the SW shearing of the Kiziba Formation it is likely that the deformation of the  
442 Kiziba Formation also occurred between 2700 Ma and 2660 Ma. The two tonalite samples  
443 collected from the TTG gneisses have similar ages (Fig. 6) with the tonalite samples that  
444 intruded the Kiziba Formation, thus, constraining the maximum age for the GSZ to ~2700 Ma.  
445 The three granite samples that intruded the TTG's and contain variably developed deformation  
446 fabrics, and were emplacement during movement on the GSZ, thus, their emplacement ages of  
447 2667 and 2680 Ma (Figs. 8 and 9) mark the main activity on the shear zone. The undeformed  
448 Nyankumbu granite constrains the minimum age movement on the GSZ at ~2640 Ma (Fig. 10).  
449 It is worth noting that the NW-trending D<sub>7</sub> shears do not appear to offset the contact of the  
450 Nyankumbu granite, indicating that shearing on the GSZ ceased before the emplacement of the  
451 Nyankumbu granite. Since, the D<sub>7</sub> shears may predate the emplacement of the 2660-2620 Ma  
452 high-K granites the activity of the GSZ can be constrained to between 2690 Ma and 2660 Ma.

453

454 *6.2. Significance of the large-scale structure for the Archean Tanzania Craton*

455

456 The GSZ has a strike length of at least 50 km (Fig. 1) and geophysical data suggest that the  
457 shear zone extends further W and E. An eastward extension of the GSZ will end up against the  
458 boundary between the Mwanza-Lake Eyasi Superterrane and the Lake Nyanza Superterrane  
459 (terminology based on Kabete et al., 2012a) while a westward extension will end up against  
460 the Proterozoic Nyakahura-Burigi Terrane. This makes the GSZ an important shear zone with  
461 large-scale consequences for the structure and tectonics of the Tanzania Craton. Assuming that  
462 the large-scale subdivision of the Tanzania Craton into a series of sub-parallel terranes  
463 separated by NW trending crustal-scale shear zones (Kabete et al., 2012a) is valid, then the  
464 GSZ will be a second order regional shear zone. However, the existence and the nature of these  
465 large-scale shear zones are yet to be confirmed by field studies and it is likely that the GSZ  
466 may represent a first order structure. It is worth noting that at least parts of the stratigraphy  
467 across the Sukumaland Greenstone Belt can be correlated. For example, the age and  
468 geochemistry of the mafic volcanics in the SGB are similar so that the Kiziba Formation north  
469 of the GSZ can be correlated with the mafic volcanics cropping out further south of the GSZ  
470 (Manya and Maboko, 2003; Manya and Maboko, 2008; Cook et al., 2016). The detrital zircon  
471 ages across the SGB and the intrusion of diorite sills suggests deposition of the Nyanzian  
472 sediments by ~2700 Ma (e.g. Borg and Krogh, 1999; Sanislav et al., 2014). The main phase of  
473 ductile deformation in the GGB is constrained by the ~ 2700 Ma maximum depositional age  
474 of the meta-volcanics, and the intrusion of the high-K granites starting at ~ 2660 Ma (Sanislav  
475 et al., 2014). The activity of the GSZ ceased at ~2660 Ma, which coincides with a switch in  
476 magmatism from a period dominated by deeply sourced magmas such as diorite and TTG to a  
477 period dominated by shallow level sourced magmas such as the high-K granites.

478

479 *6.3. The significance of the inherited cores*

480

481 Tonalite samples collected from both sides of the GSZ not only yield similar emplacement  
482 ages, but also contain similarly aged zircon xenocrysts and inherited cores. This suggests that  
483 the TTG gneisses that crop out south of the GSZ most probably extend northward beneath the  
484 GGB. The TTG gneisses south of the GSZ were initially interpreted to represent Dodoman  
485 age basement rocks on which the greenstone sequences were deposited (Bell and Dodson,  
486 1981; Gabert, 1990; Kabete et al., 2012a). However, zircon dating of a migmatitic gneiss from  
487 the southern end of the SGB returned an age on  $2680 \pm 3$  Ma (Borg and Krogh, 1999) indicating  
488 that at least some of the gneisses in the SGB are not of Dodoman age and do not constitute  
489 basement rocks to the greenstone sequences. The zircon ages presented in this study show that  
490 the age of emplacement of the TTG gneisses along the GSZ is very similar, at  $\sim 2710$  Ma,  
491 confirming previous interpretations that they do not represent Dodoman age rocks and cannot  
492 be the basement to the greenstone sequence. For example, whole rock Sm-Nd model ages from  
493 the mafic volcanics that form the base of the greenstone sequence indicate eruption at  $\sim 2820$   
494 Ma (Manya and Maboko, 2003; Manya and Maboko, 2008; Cook et al., 2016) making the  
495 deposition of the mafic volcanics at least 100 Ma older than the emplacement of the TTG's.  
496 The eruption age of the mafic volcanics is corroborated by zircon dating of interlayered rhyolite  
497 and felsic tuffs dated at  $2808 \pm 3$  Ma and  $2821 \pm 30$  Ma respectively (Borg and Krogh, 1999;  
498 Sanislav et al., 2014). It is worth noting that zircon xenocrysts and inherited cores from samples  
499 collected from both sides of the GSZ yielded ages between 2826 and 2840 Ma which are similar  
500 to the eruption age of the mafic and the interlayered felsic volcanics indicating the possibility  
501 that partial melting of the base of the greenstone sequence may have contributed to the  
502 petrogenesis of the TTG's. However, zircon cores as old as 3000 Ma or 3200 Ma found in  
503 samples SM07 and K81 cannot be correlated, at the moment, with any of the rock units from  
504 the northern half of the Tanzania Craton and raises the possibility of Dodoman age basement  
505 rocks being present at depth (Fig. 11). Kabete et al., (2012a) proposed that at least part of the

506 greenstone sequence in the SGB was deposited on rifted Dodoman basement. It is possible that  
507 >3Ga old basement rocks underlying the greenstone belt and the TTG gneisses (Fig. 11a)  
508 constitute the source of the old zircons. However, >3Ga zircon xenocrysts are found only in  
509 the proximity of the southern margin of the greenstone belt suggesting that the basement rocks  
510 do not extend further north. A second possibility is that basement rocks underlie only the TTG's  
511 (Fig. 11b). If this is the case, the GSZ must be dipping steeply N at depth to explain the  
512 presence of >3Ga zircons within tonalite that intruded the Kiziba Formation. It also implies  
513 that the GSZ has a much older history and may represent a major tectonic boundary separating  
514 two distinct terrains. The trace element and isotope geochemistry of the mafic volcanics that  
515 crop out north and south of the GSZ and form the base of the greenstone sequences in the SGB  
516 show no evidence of crustal contamination and were most probably erupted on oceanic crust  
517 (Manya and Maboko, 2003; Manya and Maboko, 2008; Cook et al., 2016). This data excludes  
518 the possibility that the greenstone sequence was deposited on old basement. However, given  
519 the fact that the geometry of the basin in which the greenstone sequence was deposited is  
520 unknown, the possibility of small fragments of rifted Dodoman age rocks incorporated in the  
521 greenstone sequence and underlying parts of the SGB (Fig. 11c) cannot be excluded.  
522 Alternatively, these 3000 to 3200 Ma zircon cores could have a detrital origin of an unknown  
523 source and were recycled through partial melting of amphibolite and sediment.

#### 524 *6.4. Crustal growth and the assembly of Tanzania Craton*

525 The zircon age data (Fig. 12) suggests that the northern part of Tanzania Craton, comprising  
526 the six granite-greenstone belts around Lake Victoria (Fig. 1), has a different age structure  
527 compared to the central part of the craton. The central part of the craton is dominated by > 3000  
528 Ma zircons (Fig. 12a) while the northern part is dominated by < 2850 Ma zircons (Fig. 12b).  
529 Indeed, all six greenstone belts (Fig. 1) from the northern part of the craton appear to have  
530 evolved within the same time period between 2600 Ma and 2850 Ma (Sanislav et al., 2014).

531 Moreover, new data suggests that some of these greenstone belts share not only a similar age  
532 history but also a similar igneous history with TTG magmatism peaking around 2700 Ma and  
533 transitioning into high-K magmatism post 2660 Ma. For example, Manya (2016a) showed that  
534 in Nzega Greenstone Belt, TTG magmatism occurred at ~ 2710 Ma and was followed by high-  
535 K magmatism at ~2670 Ma. In the SGB, the situation is almost identical, TTG magmatism  
536 occurred mainly around 2710 Ma (this study) and transitioned into high-K magmatism post  
537 2660 Ma (Sanislav et al., 2014). Musoma-Mara Greenstone Belt (Fig. 1), located further north,  
538 has a similar igneous history, although, there appears to be evidence of felsic magmatism as  
539 early as 2840 Ma (Manya et al., 2006; Sanislav et al., 2014). The central part of Tanzania  
540 Craton contains ~2815-2691 Ma greenstone fragments which, are more or less coeval with the  
541 greenstones from the northern part of the craton but they are embedded in > 3000 Ma granite  
542 and gneisses (Kabete et al., 2012b). Thus the terrane boundary between the Lake Nyanza Super  
543 Terrane and the Moyowosi-Manyoni Super Terrane (Fig. 1) could be a fundamental boundary  
544 separating two distinct cratonic nuclei. In this context the source of the zircon xenocrysts older  
545 than 3000 Ma from the TTGs along the Geita Shear Zone becomes important because may hold  
546 the key to the timing of the assembly of the Tanzania Craton. If parts of central Tanzania  
547 underlie the Sukumaland Greenstone Belt (Figs. 11a and b) and constitute the source of the >  
548 3000 Ma zircon xenocrysts than the assembly of the Tanzania Craton must have occurred prior  
549 to the emplacement of the ~2710 Ma TTGs. If that is the case dating of TTG gneisses from the  
550 northern half of the craton, particularly near the boundary with the central part of the craton  
551 should return an increasing amount of >3000 Ma zircon xenocrysts which is not the case. For  
552 example, Manya (2016a) showed that the TTG gneisses from Buzwagi mine in the Nzega  
553 Greenstone Belt (Fig. 1), located near the boundary with the central part of the craton, were  
554 emplaced at  $2713 \pm 8$  Ma, which is similar to the age of the TTG gneisses found along the Geita  
555 Shear Zone. No zircon xenocrysts were reported. Moreover, Borg and Krogh (1999) proposed

556 that prior to 2700 Ma the Nyanzian sedimentation was still active in the Sukumaland  
557 Greenstone Belt, a scenario which is at odds with the amalgamation of the craton in the same  
558 time. In fact, all shortening occurred after 2700 Ma (Sanislav et al., 2017) and is more likely  
559 that this period (<2700 Ma) coincides with the assembly of the Tanzania Craton and the Geita  
560 Shear Zone accommodated part of the strain resulted from the amalgamation of the craton.

#### 561 *6.5. Deformation style and Archean tectonics*

562 The Geita Shear Zone structural style, planar fabric with sub-horizontal stretching  
563 lineations and a linear map pattern, is similar to other Neoproterozoic shear zone systems (e.g.  
564 Dirks et al., 2002; Czarnota et al., 2010; Kabete et al., 2012a) and indicates horizontal  
565 shortening as opposed to vertical, gravity-driven, of the dome sliding deformation style typical  
566 for earlier Archean terranes (e.g. Collins et al., 1998; Van Kranendonk et al., 2007; Van  
567 Kranendonk, 2011). This type of structural style, was interpreted to result from horizontal  
568 tectonics responsible for terrane accretion in a geodynamic environment similar to modern day  
569 plate tectonics in oblique convergent settings (e.g. Percival et al., 2006; Czarnota et al., 2010;  
570 Manykiamba and Kerrich, 2012). It is worth mentioning that the structural styles typical of  
571 modern plate tectonics include accretionary mélanges, overthrust ophiolites, paired  
572 metamorphic belts and thrust and fold belts (e.g. Hamilton, 1998; Chardon et al., 1999; Stern,  
573 2005; Brown, 2007). Based on the scarcity/ lack of the forementioned examples alternative  
574 models involving subductionless continental drift were proposed to explain Archean horizontal  
575 tectonism and terrane assembly (Bédard et al., 2013). The steep nature of the Geita Shear Zone  
576 is not consistent with a thrust and fold belt scenario, even if it had a dip-slip component. This  
577 is important because as noticed by Many and Maboko (2008), the mafic volcanics of the  
578 Kiziba Formation, which form the base of the stratigraphy, are distributed throughout the  
579 Sukumaland Greenstone Belt suggesting that no large scale duplication of stratigraphy

580 occurred. That means the shortening was accommodated mainly by the localization of  
581 deformation along steeply dipping deformation zones similar to the Geita Shear Zone.

582 The fact that the shear zone is dominated by coaxial deformation and is characterized by a  
583 near horizontal mineral stretching lineation implies lateral extrusion of material (e.g. Fossen  
584 and Tikoff, 1998). The question on where the laterally extruded material ended up is a complex  
585 one and requires knowledge of the geometry of the shear zone at the time it was active. If Geita  
586 Shear Zone was linear than the angle of relative motion must change along its strike so that the  
587 coaxial component of deformation is transferred into non-coaxial deformation. Alternatively,  
588 the Geita Shear Zone had a convex shape with deformation being coaxial along the apex zone  
589 and increasingly non-coaxial along the sides. Field data suggests that to the west (Fig. 1) the  
590 shear zone preserve its linear character but it is overprinted by the Proterozoic mobile belts. To  
591 the east the entire greenstone belt changes orientation and is likely that the Geita Shear Zone  
592 curves south with possible lateral extrusion of material towards east.

## 593 **7. Conclusions**

594

595 In this contribution we provided the first field based description of a large scale shear zone  
596 from the Tanzania Craton. The Geita Shear Zone is a major shear zone that occurs in the NW  
597 part of the Tanzania Craton. The shear zone trends approximately E-W and has a strike length  
598 of at least 50 km and up to a few kilometers in width. The shear zone was active between ~2690  
599 Ma and ~2640 Ma, accommodated mostly flattening strain and most likely does not represent  
600 a terrane boundary. However, the zircon age distribution from the central and northern part of  
601 the Tanzania Craton is different, and suggests the presence of a major terrane boundary a few  
602 hundred kilometers south of the Geita Shear Zone. The deformation style of the Geita Shear  
603 Zone is similar to other Neoproterozoic shear zone systems and indicates horizontal shortening as  
604 opposed to vertical, gravity-driven deformation styles typical for earlier Archean terranes.

605

606 **Acknowledgements**

607 The authors would like to acknowledge Geita Gold Mine and AngloGold Ashanti for funding  
608 this work.

609 **Bibliography**

610 Bédard, J.H., Brouillette, P., Madore, L., Berclaz, A., 2003. Archaean cratonization  
611 and deformation in the northern Superior Province, Canada: an evaluation of plate tectonic  
612 vs vertical tectonic models. *Precambrian Research* 127, 61–87.

613 Bédard, J.H., Harris, L.B., and Thurston, P.C., 2013. The hunting of the snArc:  
614 *Precambrian Research*, 229, 20-48.

615 Bell, K., Dodson, M.H., 1981. The geochronology of the Tanzanian Shield. *Journal*  
616 *of Geology* 89, 109–228.

617 Blewett, R.S., Czarnota, K., Henson, P.A., 2010. Structural-event framework for the  
618 eastern Yilgarn Craton, Western Australia, and its implications for orogenic gold.  
619 *Precambrian Research* 183, 203–229.

620 Blewett, R.S., Shevchenko, S., Bell, B., 2004. The North Pole Dome: a non-diapiric  
621 dome in the Archaean Pilbara Craton, western Australia. *Precambrian Research* 133, 105–  
622 120.

623 Borg, G., 1992. New aspects on the lithostratigraphy and evolution of the Siga Hills,  
624 an Archaean granite-greenstone terrain in NW-Tanzania. *Zeitschrift fur Angewandte*  
625 *Geologie* 38 (2), 89-93.

626 Borg, G., Shackleton R.M., 1997. The Tanzania and NEZaire Cratons. In: de Wit,  
627 M.J., Ashwal, L.D. (Eds.) Greenstone Belts. Clarendon Press, Oxford, pp. 608-619.

628 Borg, G., Krogh, T., 1999. Isotopic age data of single zircons from the Archaean  
629 Sukumaland Greenstone Belt, Tanzania. *Journal of African Earth Sciences* 29, 301-312

630 Brown, M. 2015. Paleo- to Mesoarchean polymetamorphism in the Barberton Granite-  
631 Greenstone Belt, South Africa: Constraints from U-Pb monazite and Lu-Hf garnet  
632 geochronology on the tectonic processes that shaped the belt: discussion. *Geological Society  
633 of America Bulletin*, 127, 1550-1557.

634 Brown, M., 2007. Metamorphic conditions in orogenic belts: A record of secular  
635 change. *International Geology Reviews*, 49, 193–234

636 Card, K.D., 1990. A review of the Superior Province of the Canadian Shield, a  
637 product of Archean accretion. *Precambrian Research*, 48 , pp. 99–156

638 Cassidy, K.F., Champion, D.C., Krapež, B., Barley, M.E., Brown, S.J.A., Blewett,  
639 R.S., Groenewald, P.B., Tyler, I.M., Geological Survey of Western Australia 2006. A Revised  
640 Geological Framework for the Yilgarn Craton, Western Australia., pp. 8.

641 Chadwick, B., Vasudev, V.N. and Hegde, G.V., 2000. The Dharwar craton, southern  
642 India, interpreted as the result of Late Archaean oblique convergence. *Precambrian Research*,  
643 99, 91-111.

644 Chardon, D., Andronicos, C.L., Hollister, L.S., 1999. Largescale transpressive shear  
645 zone patterns and displacements within magmatic arcs: The coast plutonic complex, British  
646 Columbia. *Tectonics*, 18, 278–292.

647 Chardon, D., Choukroune, P., Jayananda, M., 1996. Strain patterns, decollement and  
648 incipient sagducted greenstone terrains in the Archaean Dharwar craton (south India). *Journal*  
649 *of Structural Geology* 18, 991–1004.

650 Chardon, D., Jayananda, M., Chetty, T.R.K., Peucat, J.J. 2008. Precambrian  
651 continental strain and shear zone patterns: the South Indian case. *Journal of Geophysical*  
652 *Research—Solid Earth* 113, B08402

653 Choukroune, P., Bouhallier, H., Arndt, N.T., 1995. Soft lithosphere during periods of  
654 Archaean crustal growth or crustal reworking. In: Coward, M.P., Ries, A.C. (Eds.), *Early*  
655 *Precambrian Processes*. Geological Society of London Special Publication, vol. 95, pp. 67–  
656 86.

657 Collins, W.J., Van Kranendonk, M.J., Teyssier, C., 1998. Partial convective overturn  
658 of Archaean crust in the east Pilbara Craton, Western Australia: driving mechanisms and  
659 tectonic implications. *Journal of Structural Geology* 20, 1405–1424.

660 Cook, Y.A., Sanislav, I.V., Hammerli, J., Blenkinsop, T.G., and Dirks,  
661 P.H.G.M., 2016. A primitive mantle source for the Neoproterozoic mafic rocks from the Tanzania  
662 Craton. *Geoscience Frontiers*, 7, 911-926.

663 Czarnota, K., Champion, D.C., Cassidy, K.F., Goscombe, B., Blewett, R.S., Henson,  
664 P.A., Groenewald, P.B., 2010. The geodynamics of the Eastern Goldfields Superterrane.  
665 *Precambrian Research*, 183,175–202.

666 Dirks, P. H. G. M., Charlesworth, E. G., Munyai, M. R., Wormald, R.J, 2013. Stress  
667 analyses, post-orogenic extension and 3.01 Ga gold mineralization in the Barberton  
668 Greenstone Belt, South Africa. *Precambrian Research*, 226, 157-184.

669 Dirks, P.H.G.M., Jelsma, H.A. and Hofmann, A. 2002. Accretion of an Archean  
670 greenstone belt in the Midlands of Zimbabwe, *Journal of Structural Geology*, 24, 1707-1727.

671 Fossen, H., and Tikoff, B., 1998. Extended models of transpression and transtension,  
672 and application to tectonic settings. *In: Holdsworth, R. E., Strachan, R. A. & Dewey, J. E*  
673 *(eds) 1998. Continental Transpressional and Transtensional Tectonics. Geological Society,*  
674 *London, Special Publications, 135, 15-33.*

675 Gabert, G., 1990. Lithostratigraphic and Tectonic Setting of Gold Mineralization in  
676 the Archean Cratons of Tanzania and Uganda, East Africa. *Precambrian Research* 46, 59-69.

677 Hamilton, W.B., 1998. Archean tectonics and magmatism. *International Geology*  
678 *Reviews*, 40, 1–39.

679 Hayman, P.C., Thébaud, N., Pawley, M.J., Barnes, S.J., Cas, R.A.F., Amelin, Y.,  
680 Sapkota, J., Squire, R.J., Campbell, I.H., Pegg, I., 2015. Evolution of a ~2.7 Ga large igneous  
681 province: a volcanological, geochemical and geochronological study of the Agnew  
682 Greenstone Belt, and new regional correlations for the Kalgoorlie Terrane (Yilgarn Craton,  
683 Western Australia). *Precambrian Res.* 270, 334–368.

684 Jackson, S.E., Pearson, N.J., Griffin, W.L., Belousova, E.A., 2004. The application of  
685 laser ablation-inductively coupled plasma-mass spectrometry to in situ U-Pb zircon  
686 geochronology. *Chemical Geology* 211, 47–69.

687 Jelsma, H.A., Dirks, P.H.G.M., 2002. Neo-Archaean tectonic evolution of the  
688 Zimbabwe Craton. *Geological Society, London, Special Publications* 199, 183-211.

689 Kabete, J.M., Groves, D.I., McNaughton, N.J., Mruma, A.H., 2012a. A new tectonic  
690 and temporal framework for the Tanzanian Shield: implications for gold metallogeny and  
691 undiscovered endowment. *Ore Geology Reviews* 48, 88-124.

692 Kabete, J.M., McNaughton, N.J., Groves, D.I., Mruma, A.H., 2012b. Reconnaissance  
693 SHRIMP U–Pb zircon geochronology of the Tanzania Craton: Evidence for Neoproterozoic  
694 granitoid–greenstone belts in the Central Tanzania Region and the Southern East African  
695 Orogen. *Precambrian Research* 216–219, 232–266.

696 Kloppenburg, A., White, S.H., Zegers, T.E., 2001. Structural evolution of the  
697 Warrawoona Greenstone Belt and adjoining granitoid complexes, Pilbara Craton, Australia:  
698 implications for Archean tectonic processes. *Precambrian Research* 112, 107–147.

699 Ludwig, K.R., 2003. User's Manual for Isoplot 3.00. A Geochronological Toolkit for  
700 Microsoft Excel. Berkley Geochronology Centre Special Publication No.4.

701 Manikyamba, C., and Kerrich, R., 2012. Eastern Dharwar Craton, India: Continental  
702 lithosphere growth by accretion of diverse plume and arc terranes. *Geoscience Frontiers*, 3,  
703 225-240.

704 Manya, S., 2016a. Petrogenesis and emplacement of the TTG and K-rich granites at  
705 the Buzwagi gold mine, northern Tanzania: Implications for the timing of gold  
706 mineralization. *Lithos*, 256, 26-40.

707 Manya, S., Kobayashi, K., Maboko, M.A.H., Nakamura, E., 2006. Ion microprobe  
708 zircon U–Pb dating of the late Archean metavolcanics and associated granites of the Musoma-  
709 Mara Greenstone Belt, Northeast Tanzania: implications for the geological evolution of the  
710 Tanzanian Craton. *Journal of African Earth Sciences* 45, 355–366.

711 Manya, S., Maboko, M.A.H., 2003. Dating basaltic volcanism in the Neoproterozoic  
712 Sukumaland Greenstone Belt of the Tanzania Craton using the Sm–Nd method: implications  
713 for the geological evolution of the Tanzania Craton. *Precambrian Research* 121, 35-45.

714 Many, S., Maboko, M.A.H., 2008. Geochemistry of the Neoproterozoic mafic volcanic  
715 rocks of the Geita area, NW Tanzania: implications for stratigraphical relationships in the  
716 Sukumaland greenstone belt. *Journal of African Earth Sciences* 52, 152–160.

717 Paces, J.B., Miller, J.D., 1993. Precise U–Pb ages of the Duluth Complex and related  
718 mafic intrusions, northeastern Minnesota: geochronological insights to physical,  
719 petrogenetic, paleomagnetic, and tectonomagmatic processes associated with the 1.1 Ga  
720 midcontinent rift system. *Journal of Geophysical Research* 98, 13997–14013.

721 Percival, J.A., Sanborn-Barrie, M., Skulski, T., Stott, G.M., Helmstaedt, H., and  
722 White, D.J., 2006, Tectonic evolution of the western Superior Province from NATMAP and  
723 Lithoprobe studies. *Canadian Journal of Earth Sciences*, 43, 1085–1117.

724 Polat, A., Kerrich, R., Wyman, D.A., 1998. The late Archean Schreiber–Hemlo and  
725 White River–Dayohessarah greenstone belts, Superior Province: collages of oceanic plateaus,  
726 oceanic arcs, and subduction–accretion complexes. *Tectonophysics*, 289, 295–326.

727 Quennell, A.M., McKinley, A.C.M., Aiken, W.G., 1956. Summary of the geology of  
728 Tanganyika: introduction and stratigraphy. *Tanganyika Geol. Surv. Mem.* 1 (Pt. 1) 264  
729 pp.

730 Sanislav, I. V., Brayshaw, M., Kolling, S. L., Dirks, P. H. G. M., Cook, Y. A.,  
731 Blenkinsop, T., 2017. The structural history and mineralization controls on the world-class  
732 Geita Hill gold deposit, Geita Greenstone Belt, Tanzania. *Mineralium Deposita*, 52, 257–279.

733 Sanislav, I. V., Wormald, R. J., Dirks, P. H. G. M., Blenkinsop, T. G., Salamba, L.,  
734 Joseph, D., 2014. Zircon U–Pb ages and Lu–Hf isotope systematics from late-tectonic granites,  
735 Geita greenstone belt: implications for crustal growth of the Tanzania craton. *Precambrian  
736 research*, 242, 187–204.

737 Sanislav, I.V., Kolling, S.L., Brayshaw, M., Cook, Y.A., Dirks, P.H.G.M.,  
738 Blenkinsop, T.G., Mturi, M.I., Ruhega, R., 2015. The geology of the giant Nyankanga gold  
739 deposit, Geita Greenstone Belt, Tanzania. *Ore Geology Reviews* 69, 1-16.

740 Stern, R.J., 2005. Evidence from ophiolites, blueschists, and ultrahigh-pressure  
741 metamorphic terranes that the modern episode of subduction tectonics began in  
742 Neoproterozoic time. *Geology*, 33, 557–560.

743 Van Kranendonk, M.J., 2011. Cool greenstone drips and the role of partial convective  
744 overturn in Barberton Greenstone Belt evolution. *Journal of African Earth Sciences* 60, 346–  
745 352.

746 Van Kranendonk, M.J., Collins, W.J., Hickman, A.H., Pawley, M.J., 2004. Critical  
747 tests of vertical vs. horizontal tectonic models for the Archaean East Pilbara Granite-  
748 Greenstone Terrane, Pilbara Craton, Western Australia. *Precambrian Research* 131, 173–211.

749 Van Kranendonk, M.J., Kröner, A., Hoffman, E.J., Nagel, T., and Anhaeusser, C.R.,  
750 2014. Just another drip: Reanalysis of a proposed Mesoarchean suture from the Barberton  
751 Mountain Land, South Africa: *Precambrian Research*, v. 254, p. 19–35,

752 Van Kranendonk, M.J., Smithies, R.H., Hickman, A.H., Champion, D.C., 2007.  
753 Secular tectonic evolution of Archean continental crust: interplay between horizontal and  
754 vertical processes in the formation of the Pilbara Craton, Australia. *Terra Nova*, 19, 1-38.

755 Wiedenbeck, M., Allé, P., Corfu, F., Griffin, W.L., Meier, M., Oberli, F., von Quadt,  
756 A., Roddick, J.C. and Spiegel, W., 1995. Three natural zircon standards for U-Th-Pb, Lu-Hf,  
757 trace element and REE analyses. *Geostandards Newsletter*, 19, 1-23.

758

759

760

761 **Figure captions**

762 **Figure 1.** Simplified geological map of northern half of Tanzania Craton showing the main  
763 geological units and the location of the greenstone belts (modified from Sanislav et al., 2015).  
764 Super-terrane boundaries are as proposed by Kabete et al. (2012a). SU – Sukumaland  
765 Greenstone Belt; NZ – Nzega Greenstone Belt; SM – Shynianga-Malita Greenstone Belt; IS –  
766 Iramba-Sekenke Greenstone Belt; KF – Kilimafedha Greenstone Belt; MM – Musoma-Mara  
767 Greenstone Belt.: ELVST – East Lake Victoria, MLEST- Mwanza Lake Eyasi, LNST- Lake  
768 Nyanza, MMST – Moyowosi-Manyoni, DBST – Dodoma Basement, MAST – Mbulu-Masai,  
769 NBT – Nyakahura-Burigi. Inset map of Africa showing the location of Archean blocks.

770 **Figure 2.** Geological map of Geita Greenstone Belt (modified from Sanislav et al., 2015)  
771 showing the location of dated samples. The width of the zone showing the foliation trend lines  
772 within the TTG gneisses along the southern margin of GGB marks the approximate width of  
773 the Geita Shear Zone.

774 **Figure 3.** Photographs showing outcrops of granitoids found within the Geita Shear Zone. a)  
775 mylonitic TTG near the greenstone margin; b) mylonitic TTG near the greenstone margin  
776 intruded by syn-deformation granite. Note that the granite cuts across and interfingers with the  
777 mylonitic foliation. The granite contains a similar but less intense foliation and mineral  
778 lineation with the mylonitic tonalite; c) microphotograph of altered and flattened feldspar  
779 porphyroclasts from near the contact with the Kiziba Formation. This sample was also affected  
780 by D<sub>7</sub> brittle-ductile shearing and overprinted by epidote and chlorite veins; d) symmetric,  
781 highly boudinaged quartz vein within the mylonitic TTG; e) asymmetric plagioclase

782 porphyroclasts indicating dextral movement found within the mylonitic TTG; f) TTG gneiss  
783 from ~ 400 m south of the greenstone margin having a less intense deformation fabric; g)  
784 photomicrograph showing weakly developed deformation fabric in TTG gneiss about 600 m  
785 from the greenstone margin; h and i) porphyritic granite dykes and intrusions from ~ 600 m  
786 south of the greenstone margin having only a weakly developed foliation. The feldspar  
787 phenocrysts appear undeformed; j) outcrop photograph of undeformed Nyankumbu granite that  
788 intruded and truncates the mylonitic fabric of Geita Shear Zone.

789 **Figure 4.** Stereoplots showing the poles to foliation planes and the orientation of the mineral  
790 lineations for the TTG and granites (a and b) affected by the Geita Shear Zone and for the mafic  
791 volcanics of the Kiziba Formation (c and d) that crop out near the shear zone.

792 **Figure 5.** Outcrop photographs of foliated Kiziba formation metabasalts showing an example  
793 of internal folding (a) and of asymmetrically boudinaged quartz vein (b). It is uncommon to  
794 find examples of internal folding within the metabasalts but geophysical images indicate that  
795 along the southern margin of the greenstone belt large isoclinal folds with ~E-W trending axes  
796 are present in the un-exposed areas. Asymmetrically boudinaged quartz veins indicating top to  
797 the SW movement are common.

798 **Figure 6.** Example of cathodoluminescence images of zircon grains from sample SM02 (a).  
799 Diagrams showing the  $^{207}\text{Pb}/^{206}\text{Pb}$  weighted average age (b) and the concordia plot (c) for  
800 sample SM02. Example of cathodoluminescence images of zircon grains from sample SM25  
801 (d). Note the presence of inherited cores and of high luminescence rim overgrowths. Diagrams  
802 showing the  $^{207}\text{Pb}/^{206}\text{Pb}$  weighted average age (e) and the concordia plot (c) for sample SM25.  
803 Inset diagrams show the results of analyses performed on the high luminescence rims.

804 **Figure 7.** Example of cathodoluminescence images of zircon grains from sample IH3 (a).  
805 Diagrams showing the  $^{207}\text{Pb}/^{206}\text{Pb}$  weighted average age (b) and the concordia plot (c) for

806 sample IH3. Example of cathodoluminescence images of zircon grains from sample K81 (d).  
807 Diagrams showing the  $^{207}\text{Pb}/^{206}\text{Pb}$  weighted average age (e) and the concordia plot (c) for  
808 sample K81. Note that both samples contain inherited zircon grains and rim overgrowths. Inset  
809 diagrams show the results of analyses performed on the inherited zircon cores.

810 **Figure 8.** Example of cathodoluminescence images of zircon grains from sample SM07 (a).  
811 Note the presence of inherited zircon grains and of zircon grains with complex internal pattern.  
812 Diagrams showing the  $^{207}\text{Pb}/^{206}\text{Pb}$  weighted average age (b) and the concordia plot (c) for  
813 sample SM07. Example of cathodoluminescence images of zircon grains from sample SM23  
814 (d). Note the presence of inherited cores and of high luminescence rim overgrowths. Diagrams  
815 showing the  $^{207}\text{Pb}/^{206}\text{Pb}$  weighted average age (e) and the concordia plot (c) for sample SM23.

816 **Figure 9.** Example of cathodoluminescence images of zircon grains from sample SM27 (a).  
817 Diagrams showing the  $^{207}\text{Pb}/^{206}\text{Pb}$  weighted average age (b) and the concordia plot (c) for  
818 sample SM027.

819 **Figure 10.** Example of cathodoluminescence images of zircon grains from sample SM56 (a).  
820 Diagrams showing the  $^{207}\text{Pb}/^{206}\text{Pb}$  weighted average ages (b) and the concordia plot (c) for  
821 different zircon populations from sample SM56. Example of cathodoluminescence images of  
822 zircon grains from sample SM57 (d). Diagrams showing the  $^{207}\text{Pb}/^{206}\text{Pb}$  weighted average age  
823 (e) and the concordia plot (c) for the youngest zircon population for sample SM57. Both  
824 samples contain inherited zircon grains having similar ages with the surrounding TTG and  
825 granites.

826 **Figure 11.** Series of cartoons illustrating three possible scenarios involving the presence at  
827 depth of > 3 Ga basement rocks to explain the old zircon ages found in the TTG and granite  
828 samples from both sides of the Geita Shear Zone. In all three scenarios, the melt gets  
829 contaminated with old zircons during ascent. It is worth noting that, so far, old zircon grains

830 were found only within the vicinity of the Geita Shear Zone. a) Old basement underlying both  
831 the TTG and the greenstone. If that is the case old zircon grains could be found further south.  
832 b) Old basement underlying only the TTG and GSZ dips north at lower depths. If this is the  
833 case, GSZ could be a major tectonic boundary separating two distinct terrains. It would also  
834 suggest that the shear zones initiated before 2700 Ma. Old zircon grains could be found further  
835 south. c) A fragment of rifted old basement underlies part of the greenstone belt. If that is the  
836 case old zircon grains would occur only in certain domains.

837 **Figure 12.** Diagrams showing the distribution of zircon ages for central Tanzania (a) and for  
838 northern Tanzania (b). Data from Sanislav et al., (2014) updated with new age data from Many  
839 (2016a) and present study.

840

841